

FINAL REPORT

to

Dr. Mirsajadi Hassan
Watershed Assessment Section, Division of Water Resources,
Delaware Natural Resources and Environmental Control (DNREC)
89 Kings Hwy
Dover, DE 19901
Email: Hassan.Mirsajadi@state.de.us

**Vertical Profiles of Radioisotopes, Nutrients and Diatoms in Sediment Cores
from the Tidal Murderkill River Basin:
A Historical Analysis of Ecological Change and Sediment Accretion**

PCER Report No. 10-01

By
Drs. David Velinsky, Christopher Sommerfield¹ and Don Charles
Patrick Center for Environmental Research
The Academy of Natural Sciences
Philadelphia, PA 19103

¹University of Delaware
College of Earth, Ocean and Environment
School of Marine Science and Policy
Lewes, DE

June 29, 2010

TABLE OF CONTENTS

	Page
List of Tables	iii
List of Figures	iv
Executive Summary	v
 A	
Introduction	1
A1 Background	1
A2 Objectives of Study	3
A3 Study Area	3
 B	
Field and Laboratory Methods	5
B1 Field Sampling	5
B2 Laboratory Methods.....	5
B2.1 Radioisotope Measurements and Sedimentation Rates	6
B2.2 Total Organic Carbon and Nitrogen.....	8
B2.3 Total Phosphorus.....	8
B2.4 Stable Isotopes of Carbon and Nitrogen	8
B2.5 Diatoms	9
 C	
Results and Discussion	10
C1 Sediment Properties and Sediment Accumulation.....	10
C1.1 Sediment Properties.....	10
C1.2 Accretion and Sediment Accumulation Rates.....	10
C1.3 Radionuclide Inventories and Focusing Factors	12
C2 Nutrients/Eutrophication.....	14
C2.1 Sediment Total Carbon, Total Nitrogen and Total Phosphorus	14
C2.2 Stable Isotopes of Carbon and Nitrogen	15
C2.3 Diatom Analysis and Assemblages	17
C3 Historical Analysis.....	21
C3.1 Changes in Nutrients and Diatoms in Murderkill River Cores	22
C3.2 Changes in Marsh Ecology	25
C3.3 Nitrogen and Phosphorus Accumulation Rates.....	26

TABLE OF CONTENTS (cont)

	Page
D	
Summary and Conclusions	28
E	
Acknowledgments	30
F	
References	31
G	
Tables	39
H	
Figures	58
I	
Appendices	77
Appendix I: Data Tables and QA Documentation (electronic only).....	I

LIST OF TABLES

Table 1: Core locations and collection dates	40
Table 2: Bulk sediment properties and radioisotope data for core MK-1.....	41
Table 3: Bulk sediment properties and radioisotope data for core MK-2.....	43
Table 4: Bulk sediment properties and radioisotope data for core MK-3.....	45
Table 5: Bulk sediment properties and radioisotope data for core MK-4.....	47
Table 6: Summary data for radioisotope analysis and dating.....	49
Table 7: Concentrations of various parameters for core MK-1	50
Table 8: Concentrations of various parameters for core MK-2	51
Table 9: Concentrations of various parameters for core MK-3	52
Table 10: Concentrations of various parameters for core MK-4	53
Table 11: Diatom metrics for core MK-1	54
Table 12: Diatom metrics for core MK-2	55
Table 13: Diatom metrics for core MK-3	56
Table 14: Diatom metrics for core MK-4	57

LIST OF FIGURES

Figure 1: Generalized schematic of nitrogen and phosphorus cycling in wetlands.....	59
Figure 2: Study area and core locations.....	60
Figure 3: Tripod/pulley system in the Murderkill River.....	61
Figure 4: Geochronology for core MK-1.....	62
Figure 5: Geochronology for core MK-2.....	62
Figure 6: Geochronology for core MK-3.....	63
Figure 7: Geochronology for core MK-4.....	63
Figure 8: Sediment organic carbon, C/N and sediment phosphorus distribution with depth	64
Figure 9: Depth distribution of the isotopic composition sediment N ($\delta^{15}\text{N}$) and C ($\delta^{13}\text{C}$)	65
Figure 10: Distribution of diatom indicator species with depth: MK-1.....	66
Figure 11: Distribution of diatom indicator species with depth: MK-2.....	67
Figure 12: Distribution of diatom indicator species with depth: MK-3.....	68
Figure 13: Distribution of diatom indicator species with depth: MK-4.....	69
Figure 14: Diatom metrics	70
Figure 15: Relationship between total sediment N and diatom metrics	71
Figure 16: Relationship between the accumulation rate of sediment N and P and the diatom metric for eutrophic species in each core.....	72
Figure 17: Concentrations of total sediment nitrogen (TN) and the nitrogen isotopic composition of TN ($\delta^{15}\text{N-TN}$) from 1860 to 2008	73
Figure 18: Concentrations of total sediment phosphorus (TP) and the carbon isotopic composition of total carbon ($\delta^{13}\text{C-TC}$) from 1860 to 2008	74
Figure 19: Diatom metric for eutrophic species from 1860 to 2008.....	75
Figure 20: Nitrogen and phosphorus accumulation with time.....	76

Executive Summary

This study involved the chemical and biological analysis of four sediment cores from the tidal Murderkill River in Kent County, DE. These cores were collected from upstream near Frederica to downstream (near Bowers Beach) and were dated using ^{210}Pb and ^{137}Cs radiometric methods. All cores provided sufficient temporal coverage (> 80-100 yrs) for detailed chemical analysis. *The objective of this study was to obtain a historical perspective on nutrient inputs and ecological response into the Murderkill River using dated sediment cores, and to evaluate whether pollution controls have been effective.* We evaluated historical trends in nutrients (i.e., sediment phosphorus and nitrogen) and assessed if the historical record of eutrophication could be derived from algal analysis (i.e., diatom species in the cores) along with other indicators of potential ecosystem change (e.g., stable isotopes of carbon ($\delta^{13}\text{C-OM}$) and nitrogen, ($\delta^{15}\text{N-TN}$)).

The range and average sediment accumulation rates derived from the ^{210}Pb and ^{137}Cs data are similar to other areas within the Delaware Estuary and range from 0.33 to 0.74 cm/yr. There was good agreement between the ^{137}Cs rates and ^{210}Pb using the constant initial concentration model.

Sediment organic carbon (SOC) concentrations for the four cores ranged between 2.3% and 34.3% on a dry weight basis (dw) with an average of $13.4 \pm 8.1\%$ SOC ($\pm 1\sigma$). Similarly, total nitrogen (TN) ranged from 0.21 to 1.7% N with an overall average of $0.68 \pm 0.37\%$; whereas total sediment phosphorus (TSP) ranged from 0.03 to 0.21% TSP with an overall average of $0.07 \pm 0.03\%$. Similar concentrations were found for TN and TSP in marsh cores taken from the tidal river in the summer of 2007 (Chesapeake Biogeochemical Associates, 2007). Sediment nitrogen increased in only MK-2 whereas TSP showed only a small increase towards the core top. Sediment N and P accumulation rates increased upward in the sediment column starting around the mid to late 1970s. This is approximately the same time frame as the onset of effluent discharge from the Kent County Wasted Water Treatment Plant (KC WWTP).

Analysis of the diatom assemblages and metrics indicate a shift toward more eutrophic species starting in the late 1940-1950s (cores MK-1 and MK-2) or 1970s (core MK-3), but no clear trend was observed in core MK-4. Although no significant relationship was found between the eutrophic diatom metrics and the concentration and accumulation rates of nitrogen and phosphorus in the cores, additional statistical analyses are necessary to identify specific trends. For example, cross correlations should be used to account for the lag in diatom response to nutrient changes, or for processes related to sediment nutrient re-mobilization. Also, more

analyses can be conducted on select groups of diatom species known to have stronger relationships with nutrient concentration. The study of sediment cores from Murderkill tidal river revealed an important aspect of the assemblages that affect our ability to reconstruct environmental changes. The diatom assemblages included many brackish or marine species and little is known about their nutrient and habitat preference; further investigations with simultaneous water quality measurements and quantification of diatom species from coastal environments are necessary to better estimate their ecological requirements and then to understand historical changes with sediment cores.

An interesting finding of this study was the shift of diatom flora from freshwater to brackish-marine species in cores MK-1 and MK-2, along with a change in the stable isotopic composition of sediment organic carbon. The organic carbon shifted from more negative values to more positive values indicating a change from C3 plants to C4 plants with time. These data suggest a change of salinity in the upper tidal reaches and that these changes shifted the marsh to its current composition in a time frame of a few decades. It is unclear as to what caused this change but it is thought that some hydrological modification occurred that allowed higher salinity water to move upstream. To validate this change, further analysis of sediment proxies need to be accomplished, such as benthic foraminifera composition, bromide (an indicator of salinity) of organic matter and additional cores. The data could suggest a potential change due to sea level rise, salinity intrusion, and changes in marsh ecology. The potential influence of such factors needs to be explored by analyzing relationships between climate and sea level records and the core sediment proxies.

Lastly, N and P accumulation rates over time have increased two fold since approximately 1975-1980 in a similar time frame as loadings from the KC WWTP. Preliminary estimates of marsh burial suggest that current burial rates can remove a substantial fraction of both N and P; however, recycling rates need to be considered for more accurate estimates. This study provides important data and results that can be incorporated into an ecosystem model for the tidal river to help determine changes in water quality.

A) Introduction

A1: Background

Nutrients, trace metals and organic contaminants in water are derived from many sources. Natural sources of metals include the weathering products of soils which are then transported in the dissolved or particulate phases. Anthropogenic sources of metals and organic contaminants are introduced to the water via atmospheric deposition, industrial discharges (e.g., mining, metal processing, manufacturing), municipal discharges (waste water treatment), and stormwater runoff of contaminated parcels. Many of these same sources are also noted for discharges of nitrogen and phosphorus, especially waste water treatment facilities and agricultural runoff (via surface runoff and indirectly through groundwater flows). Due to the particle-reactive nature of most trace metals, phosphorus, and organic compounds, sediments are potential repositories for contaminants and, under certain conditions, can be used to provide a historical record of pollution (Simpson et al., 1983; Orson et al., 1990; 1992; Valette-Silver, 1993; Hornberger et al., 1999; Cooper and Brush, 1993; Church et al., 2006; Ridgeway and Shimmiel, 2002; Velinsky et al., 2007; Hartzell et al., 2010, and others). With minimal post-depositional diagenetic remobilization, chemical transformations, biological mixing, and hydraulic processes, the sediment column can record the chronology of nutrient and chemical contaminant loading and burial in estuarine waters.

Coring information can be used for the construction of sediment budgets (Schubel and Hirschberg, 1977; Brush et al., 1982; Officer et al., 1984) and to understand chemical/nutrient accumulation in aquatic environments (Owens and Cornwell, 1995; Cornwell et al., 1996; Latimer and Quinn, 1996; Van Metre and Callender, 1997; Church et al., 2006; Velinsky et al., 2007; 2010; Hartzell et al., 2010). Also, dated sediment cores can provide loadings information of contaminants or chemicals over time (e.g., Rippey and Anderson, 1996; Santschi et al., 2001). However, specific chemicals, such as nitrogen and phosphorus can undergo substantial diagenetic re-mobilization depending on the redox conditions (see Burdige, 2006) and as such accumulation records derived from sediment cores maybe limited in certain environments.

Phosphorus is transported to the oceans from rivers and estuaries and up to 90% is in a particulate form (Meybeck, 1982; Froelich et al., 1982; Lebo, 1991; Follmi, 1996; Litke, 1999; Jordan et al., 2008). Once bound phosphorus is buried, it has the potential to be recycled back into the water column depending on many factors (Froelich et al., 1982; Boynton et al., 1995;

Jordan et al., 2008; **Figure 1**). In some sub-tidal environments, only a small portion of the buried phosphorus (e.g., 5 to 30%) is retained, hindering interpretation of anthropogenic sources with time. However, Kahn and Brush (1994), Cornwell et al. (1996), and Church et al. (2006) found no major change in phosphorus with depth in sediment cores, and concentration profiles tracked changes in loadings over time. Controlling factors that would enhance retention of buried phosphorus are the magnitude and rate of phosphorus loadings, sediment accumulation rate, and the magnitude and rate of remobilization of phosphorus in the sediments.

Nitrogen is transported in rivers and estuaries mostly in the dissolved form and its cycling is much more complex (Meybeck, 1982, Van Breemen et al. 2002; Castro et al., 2003; **Figure 1**). As such, retention of nitrogen in coastal and riverine sediments is potentially more restricted than for phosphorus. As with phosphorus, controlling factors may include the rate of sediment accumulation and the magnitude of nitrogen loading. The stable isotopes of nitrogen ($^{14}\text{N}/^{15}\text{N}$) can help determine the source, fate and cycling of nitrogen in a water body and could be reflected in the nitrogen buried in sediments (see Kendall, 1998). For example, Church et al. (2006) showed an increase in $\delta^{15}\text{N}$ of sediment N with time that is reflective of increases in urbanization in the Delaware Estuary, similar to other urbanized waters (McClelland et al., 1997; Kendall, 1998; Lake et al., 2001; Ulseth and Hershey, 2005).

Sediment core records for rivers and estuaries can be extremely useful in determining if pollution control actions were/are effective in reducing contaminant loadings, as well as providing a time frame for system response (Owens and Cornwell, 1995; Zhang et al., 1993; Smol, 2008). In addition, biological material, and in some cases chemical information, retained in the sediments can help determine changes in ecosystem processes and health (Smol, 2008; Potapova and Charles, 2007; Bennion et al., 2001; Cooper, 1999; 1995; Cooper and Brush, 1993). Specific diatoms and foraminifera can be used as proxies to determine changes in response to watershed and estuarine changes in nutrient loadings and concentrations. Pollen content of sediments has been very useful in describing past changes in vegetation in related to land use changes over time (Cooper and Brush, 1993; Brush and Brush, 1994), whereas biogenic silica, a chemically determined proxy, in conjunction with other chemical proxies can be used to determine aquatic productivity (see Conley and Schelske, 2001). Overall, chemical and biological proxies in sediments cores can provide a wealth of information about watershed and

aquatic ecosystem change and sediment core data is especially important for modeling programs in which forecasts are developed by extending past trajectories.

A2: Objectives of Study

The objective of this study was to collect sediment cores from the tidal and estuarine region of the Murderkill River and determine the chronology of carbon, nitrogen and phosphorus deposition, their loading histories, and related ecological responses. To meet this objective we obtained and analyzed the chemical characteristics of sediment cores from marsh depositional areas fringing the estuary. In addition to providing insight into the temporal variability of nitrogen and phosphorus in the estuary, we also sought to understand how the trophic status of the system has changed over decadal time scales.

A3: Study Area

The Murderkill River Basin occupies 275 km² in the southeastern portion of Kent County, Delaware. The Murderkill River is the main branch of the watershed system and flows approximately 37 km from its headwaters near Felton, Delaware, to its confluence with the Delaware Bay at Bowers Beach (**Figure 2**). The lower portion from upstream of the Route 113 Bridge at Frederica, Delaware, to the mouth is tidally influenced and is approximately 5 km in length. The current land use of the watershed is agriculture (55%), wooded/forest (17%), wetlands (6-9%), urban areas (14%) and open water (<2%) (DNREC, 2005).

Changes in the Murderkill River Watershed and Nutrient Enrichment

The water quality of the Murderkill is affected by persistent pollution impacts (i.e., eutrophication and low dissolved oxygen) from agricultural runoff and wastewater discharges, in addition to having somewhat restricted tidal flushing (Aurand and Daiber, 1973; deWitt and Daiber, 1974; DNREC, 2005). Nutrient over-enrichment and low dissolved oxygen are two environmental issues of concern in the Murderkill watershed (DNREC, 2007). Major sources are point source discharges, surface runoff from agricultural fields, animal-raising operations, septic tanks, and historic uses as noted in elevated nitrogen in groundwater discharges. In the mid to late 1970s, a large wastewater treatment plant was constructed in Frederica which, at the time, had a planned mean daily discharge of $38 \times 10^3 \text{ m}^3$ (de Michele, 1972 as cited in deWitt

and Daiber, 1974). Currently, there is one major discharger (Kent County Facility, DE0020338) and two minor (Harrington STP and Canterbury Crossing MHP) to the river. The Kent County facility discharges (~ 11 MGD; $\sim 42 \times 10^3 \text{ m}^3/\text{day}$) directly to the tidal portion of the Murderkill River. Ullman et al. (2010) estimate 2007 nutrient loads from the Kent County facility based on discharge monitoring reports. Total external (new) loads of N and P to the tidal portion vary over the year. Nitrogen loads in 2007 ranged from 20 to 110 metric tons per month and were mainly derived from upstream watershed inputs with lesser amounts from atmospheric deposition and the Kent County Waste Water Treatment Plant (KC WWTP). Phosphorus loads to the tidal waters ranged from 0.5 to 8 metric tons per month in 2007 with $>\sim 80\%$ derived from the Kent County WWTP and lesser amounts from the watershed and atmospheric deposition.

Historically, the tidal waters of the sub-estuary are considered to have abnormally low rates of dissolved oxygen (deWitt and Daiber, 1974) of between 63 to $\sim 90 \mu\text{M O}_2$ during the summertime. In addition, Aurand and Daiber (1973) measured substantial concentrations of dissolved nitrate of between 40 to $100 \mu\text{M N}$ in winter decreasing to 5 to $20 \mu\text{M}$ in the summer. This is most likely the results of agricultural and animal-raising operations. Recent concentrations of dissolved inorganic N (DIN) ranged from 100 to $300 \mu\text{M N}$ in the tidal river (fall, 2008) and, for dissolved inorganic P (DIP), from 2 to $8 \mu\text{M P}$. The river is currently listed on Delaware's 303(d) list of impaired waters (DNREC, 2005; 2007) due to high bacteria levels, low dissolved oxygen and excessive nutrients. Based on modeling analyses, both point and non-point sources of nitrogen, phosphorus, and organic material (i.e., BOD) must be reduced to meet Delaware water quality standards and targets. In 2005, EPA and DNREC implemented a TMDL for excess nutrient inputs to the river (high and low flow conditions). Monitoring shows that 310-450 kg of N and 130-153 kg of P are discharged daily into the river (US EPA, 2005) from the Kent County facility. The waste load allocation for the facility, set by a TMDL, was 340 kg N/day and 28 kg P/day (reduction of 80%) with compliance gauged on a monthly basis (DNREC, 2006).

In general, there is a lack of historical water monitoring data, i.e., from the 1940s/1960s to the present, to assess the changes in land use and nutrient addition on the water quality (e.g., dissolved oxygen, plant ecology, marsh dynamics) of the tidal river. The goal of this project was to obtain a historical perspective on nutrient inputs and ecological response into the Murderkill River using dated sediment cores, and to evaluate whether pollution controls have been effective.

B) Field and Laboratory Methods

B1: Field Sampling

Sediment cores were collected on October 10, 2008, by staff from The Academy of Natural Sciences and University of Delaware (UDEL) at four locations in the tidal river and estuary (**Figure 2; Table 1**). A total of eight cores were obtained, and all provided good sediment chronologies based on ^{210}Pb and ^{137}Cs measurements.

Cores were collected on the marsh surface during mid to low tide. Core locations were from the interior of the marsh, away from any obvious disturbances (e.g., creek banks and ditching). At each site two cores were obtained, one for chemical analysis and the other for stratigraphic descriptions of each site. Push-piston cores of approximately 1 to 1.5 m in length were retrieved by a tripod/pulley system (**Figure 3**). The cores were taken to the laboratory, extruded vertically, and sectioned in 2-cm intervals. Samples were stored in pre-cleaned jars at -10°C at either Academy or UDEL facilities. Chain-of-custody procedures were followed from the time of collection, shipping and until the analyses were completed.

B2: Laboratory Methods

Published laboratory clean-techniques were used throughout (Church et al., 2006; US EPA, 1997; APHA, AWWA and WEF, 1995) using protocols as outlined in standard operating procedures (SOPS) at the Academy of Natural Sciences and University of Delaware. All materials coming in contact with the samples were either glass or metal and were cleaned of any contaminants prior to use. Sample ID forms were used and each sample was given a unique laboratory number for sample tracking.

Sediments were analyzed for the following parameters at laboratories operated by the Academy of Natural Sciences (Patrick Center): organic carbon, total nitrogen and total phosphorus, and stable isotopes of carbon and nitrogen. In addition, specific sections were analyzed for diatoms via sample digestion, mounting and glass slide light microscopy. Sediments for radioisotope measurements (^{210}Pb and ^{137}Cs) were analyzed at University of Delaware (School of Marine Science and Policy). Below are brief descriptions of each chemical, biological, or physical method.

B2.1: Radioisotope Measurements and Sedimentation Rates

Sediment bulk density (ρ_d) and loss-on-ignition (LOI) measurements were made for each core section to aid interpretation of the downcore radionuclide and chemical data. Dry-bulk density was calculated from porosity (ϕ) using representative values of interstitial fluid density (ρ_f) and mineral density (ρ_m) according to:

$$\rho_d = (1 - \phi) \rho_m \quad (1)$$

where $\rho_m = 2.65 \text{ g/cm}^3$. Porosity was computed gravimetrically from water content (W_c) and using an assumed pore water density of $\rho_f = 1.0 \text{ g/cm}^3$:

$$\phi = \left(\frac{W_c \rho_m}{W_c \rho_m + (1 - W_c) \rho_f} \right) \quad (2)$$

where $W_c = W_{wet} - W_{dry} / W_{wet}$. Dry sediment weight was determined by drying the sectioned wet sediment in a convection oven at 100°C for 24 h. The dried sediment was then ground to a fine powder for LOI determination and radionuclide analysis.

LOI measurements were used to quantify the relative proportion of organogenic (combustible) and minerogenic (residual ash) materials in the sediments. Following the method of Heiri et al. (2001), a 4-g quantity of sample powder was combusted at 550°C in a muffle furnace for 4 h. LOI was computed as follows:

$$LOI = \frac{W_{dry} - W_{ash}}{W_{dry}} \cdot 100\% \quad (3)$$

where W_{dry} is the weight of sample previously dried at 100°C , and W_{ash} is the weight of the residual ash.

To develop sediment chronologies for the sediment column, measurements of ^{210}Pb ($t_{1/2} = 22.3$ years) and ^{137}Cs ($t_{1/2} = 30.1$ years) were made by gamma spectroscopy of the 46.5 and 661.6 keV photopeaks, respectively (Cutshall et al., 1983; Wallbrink et al., 2002). Lead-210 is a natural radionuclide of the U-238 decay series, whereas ^{137}Cs was introduced to aquatic waters via nuclear weapons testing fallout in ca. 1954. Both radionuclides are delivered to the marsh surface through a combination of direct atmospheric deposition and lateral transport by tidal flooding. Powder samples for radionuclide analysis were placed in a 60-ml plastic jar and counted for 24-48 h on a Canberra Model 2020 low-energy Germanium detector (LEGe). The

concentration (activity) of excess ^{210}Pb was determined by subtracting the activity of its parent nuclide ^{214}Bi (609.3 keV) from the total activity of ^{210}Pb ($^{210}\text{Pb}_{\text{ex}} = ^{210}\text{Pb}_{\text{tot}} - ^{214}\text{Bi}$), assuming secular equilibrium between ^{214}Bi and ^{210}Pb . Detector efficiencies were determined from counts of NIST Standard Reference Material 4357 (Inn et al., 2001). The counting geometry of the samples was kept identical to that of the NIST standard such that a self-absorption correction for ^{210}Pb was not necessary. Confidence limits reported with radioisotope data are the propagated one-sigma background, calibration, and counting errors.

Two approaches were used to reconstruct the accumulation history of the marsh sediment column recovered in cores. The first is based on the downcore activity profile of ^{210}Pb , and the second uses the depth distribution of ^{137}Cs activity. For the first approach, an accretion rate (cm/yr) for each coring site was determined from the excess ^{210}Pb profile following the Constant Initial Concentration (CIC) model (Robbins, 1978). This model assumes that the specific activity (dpm/g) of excess ^{210}Pb deposited on the marsh surface remains constant through time (steady state). In other words, variations in mineral deposition rate do not alter the initial concentration of excess ^{210}Pb . At steady state, excess ^{210}Pb decreases exponentially with depth in the sediment column following:

$$A = A_0 \exp(-\lambda z/S) \quad (4)$$

where A is the excess activity of ^{210}Pb at depth z , A_0 is the initial activity of excess ^{210}Pb , λ is the decay constant for ^{210}Pb (0.03114 years), and S is the vertical accretion rate. On a plot of $\ln A$ versus z the slope of a linear regression line is proportional to the accretion rate. In the CIC model the age of a sediment interval relative to the year of sampling is derived from a single value of S averaged over the entire ^{210}Pb profile. At the marsh sites sampled in this study the ^{210}Pb method provided approximately 80–100 years worth of accumulation history. The mass accumulation rate (mass/area/time) of the sediment column was determined by plotting A as a function of cumulative mass, the product of ρ_d and z .

The second chronological method is based on the activity peak of ^{137}Cs preserved in the sediment column as an indicator of ca. 1964, the year of maximal fallout of ^{137}Cs from the atmosphere (Ritchie and McHenry, 1990). Dividing the sediment depth of the ^{137}Cs activity peak by the time period between ca. 1964 and the year of core collection gives an accretion rate averaged over the past ~44 years. The corresponding mass accumulation rate is determined by

plotting ^{137}Cs activity against cumulative mass. The advantage of ^{137}Cs chronology over the ^{210}Pb method is that it more closely provides the absolute age of sediments deposited after ca. 1964.

Sediment inventories of excess ^{210}Pb and ^{137}Cs were computed to compare the relative amount of radionuclide deposited among the four coring sites (MK-1 through MK-4). Inventories were computed as:

$$I_o = \sum_i \rho_{di} x_i A_i \quad (5)$$

where ρ_d is the dry-bulk density, x is the sediment thickness, A is the radionuclide activity, and the i operator indicates the i th depth interval.

The following conditions are implicit in all ^{210}Pb and ^{137}Cs dating methods: 1) mixing by burrowing organism has not augmented burial of particles and adsorbed radionuclide; 2) the radionuclide is chemically immobile in the sediment column; and 3) the sedimentary record is complete and not punctuated by prolonged episodes of non-deposition or erosion. As elaborated later, these assumptions appear to have been met in this study for all cores.

B2.2: Total Organic Carbon and Total Nitrogen

Total organic carbon and total nitrogen were measured using a CE Flash Elemental Analyzer following the guidelines in EPA 440.0, manufacturer instructions and ANSP-PC SOP. Samples were pre-treated with acid to remove inorganic carbon.

B2.3: Total Phosphorus

Total sediment phosphorus was determined using a dry oxidation method modified from Aspila et al. (1976) and Ruttenberg (1992). Solubilized inorganic phosphorus was measured with standard phosphate procedures using an Alpkem Rapid Flow Analyzer. Standard reference material (spinach leaves) and procedural blanks were analyzed periodically during this study. All concentrations were reported on a dry weight basis.

B2.4: Stable Isotopes of Carbon and Nitrogen

The stable isotopic composition of sediments was analyzed using a Finnigan Delta XL coupled to an NA2500 Elemental Analyzer (EA-IRMS). Samples were run in duplicate or triplicate with the results reported in the standard δ (‰) notation: $\delta X = (R_{\text{sample}}/R_{\text{standard}}) - 1$ X

1000; where X is either ^{13}C or ^{15}N and R is either $^{13}\text{C}/^{12}\text{C}$ or $^{15}\text{N}/^{14}\text{N}$. The $\delta^{15}\text{N}$ standard was air ($\delta^{15}\text{N} = 0$), and for $\delta^{13}\text{C}$ the standard is the Vienna PeeDee Belemite (VPDB) limestone that has been assigned a value of 0.0 ‰. Analytical accuracy was based on the standardization of the UHP N_2 and CO_2 used for continuous flow-IRMS with IAEA N-1 and N-2 for nitrogen and IAEA sucrose for carbon, respectively. An in-house calibrated sediment standard was analyzed every tenth sample. Generally, precision based on replicate sample analysis was better than 0.2‰ for carbon and 0.6‰ for nitrogen.

B2.5: Diatoms

About 1-g core sediment was subsampled and the organic component was oxidized with 70% nitric acid while heated in a CEM microwave (165°C) for 1.5 h. Diatoms were repeatedly allowed to settle for 24 hours and the supernatant was decanted until it reached a neutral pH. A measured amount of digested sample was dripped onto a microscope cover slip and dried. Cover slips were then mounted onto slides using a high refractive index mounting medium (Naphrax™). Diatoms were counted and identified using a Leica DM LB2 microscope equipped with DIC optics. At least four hundred valves were counted for each slide at 1000x magnification, unless diatom concentration was too low to allow a 400-valve count. More details on standard Phycology Section operating procedures for diatom analysis can be found in “Protocols for the analysis of algal samples collected as part of the USGS National Water Quality Assessment Program” (Charles et al., 2002; <http://diatom.ansp.org/nawqa/protocols.asp>). Diatom species identifications were made using the extensive diatom library at ANSP (Charles et al., 2002). Several diatom community metrics were calculated based on species autecological preferences originally assembled by van Dam et al. (1994). Metrics were calculated using the Phyco-Aide program developed at ANSP. Assemblages in each core were analyzed using Principal Component Analysis (PCA). The scores of each taxon along PCA axis 1 were used to order the sequence of taxa in diatom stratigraphic profiles.

C) Results and Discussion

C1: Sediment Bulk Properties and Sediment Accumulation

C1.1: Sediment Bulk Properties

The cored deposits consisted of clayey silt with variable quantities of living and dead plant material. Dry-bulk densities ranged from 0.11 to 0.76 g/cm³ overall, and each core displayed a slightly different depth profile (**Figures 4-7; Tables 2-5**). In a compositionally uniform sediment column, bulk density generally increases with depth with compaction and expulsion of porewater. However, this general trend was not evident in all cores presumably on account of down-core variations in organic matter and grain size. In cores MK-1 and MK-2, bulk density initially increased with depth to ~40 cm but then decreased toward the core bottom. Bulk density in core MK-3 was more-or-less uniform with depth to 80 cm, increasing sharply to the bottom at 100 cm. Among cores bulk density was most variable in depth in MK-4, ranging from 0.3 to 0.76 g/cm³ with no net depth trend.

Based on LOI measurements the organic content of cores ranged from 9.6 to 71% by weight (**Figures 4-7; Tables 2-5**). Organic content was highest in cores from the upriver sites (MK-1 and MK-2), particularly in the lower the sediment column where it reached ~70%. In cores MK-1 and MK-2, organic content increased downcore abruptly around 42–46 cm sediment depths, whereas in cores MK-3 and MK-4 it decreased only gradually from top to bottom. LOI varied inversely with dry-bulk density overall, suggesting that the values of bulk density measured in cores are partly a function of the concentration of organogenic material. In summary, the sediment properties data suggest that there was a major change in the nature of material deposited at sites MK-1 and MK-2 over time. Other chemical and biological indicators reflect changes as well (see below).

C1.2: Accretion and Sediment Accumulation Rates

The profile of excess ²¹⁰Pb activity for core MK-1 exhibited a monotonic decrease downcore to a depth of 58 cm, indicative of steady-state sediment accumulation and radioactive decay (**Figure 4**). The measured accretion rate is 0.74 cm/yr, and the corresponding mass accumulation rate is 0.20 g/cm²/yr. Activities of ¹³⁷Cs increased upcore from the depth of first

occurrence at 44-48 cm to a sharp peak at 28-32 cm, above which activities decreased to lower, but detectable, values near the core top (**Figure 4; Table 2**). The shape of the ^{137}Cs profile for this core (and for the other cores in the study area) is broadly consistent with the record of ^{137}Cs atmospheric fallout in U.S. Mid-Atlantic region since 1954 (Olsen et al., 1981). Accretion and mass accumulation rates calculated from the depth of the ^{137}Cs activity peak are 0.71 cm/yr and 0.19 g/cm²/yr, respectively.

Core MK-2 exhibited ^{137}Cs and ^{210}Pb distributions similar to those observed in core MK-1 (**Figure 5; Table 3**). Excess ^{210}Pb activity extended to 90 cm, and rates of accretion and accumulation calculated from the decay profile are 0.74 cm/yr and 0.13 g/cm²/yr. ^{137}Cs activities increased upcore from the depth of first occurrence at 44-48 cm to a distinct peak centered at 28-32 cm, and then decreased to near-zero activities at the core top. Although ^{137}Cs was detectable in the 44-48 cm sediment interval, the activity was just above the minimum detectable activity. Accretion and mass accumulation rates based on the ^{137}Cs peak are 0.71 cm/yr and 0.12 g/cm²/yr, respectively.

Excess ^{210}Pb activity in core MK-3 extended to a depth of 50 cm, and the decay profile indicates an accretion and mass accumulation rate of 0.60 cm/yr and 0.14 g/cm²/yr, respectively (**Figure 6; Table 4**). ^{137}Cs activities increased upcore from first occurrence at 32-36 cm to a distinct peak centered at 16-20 cm and thereafter decreased to near-zero at the core top. Accretion and mass accumulation rates based on the depth of the ^{137}Cs peak were 0.44 cm/y and 0.10 g/cm²/yr, respectively.

As in core MK-3, excess ^{210}Pb activity in core MK-4 extended to a depth of 50 cm but with more scatter about the trend line at depth. Accretion and mass accumulation rates calculated from the ^{210}Pb decay profile are 0.33 cm/yr and 0.17 g/cm²/yr, respectively. The first occurrence of ^{137}Cs activity in core MK-4 was centered at 20-22 cm, with a broad peak present at 8-16 cm (**Figure 7; Table 5**). The ^{137}Cs accretion and mass accumulation rates for this site were 0.31 cm/yr and 0.16 g/cm²/yr, respectively, which were the lowest rates among the four sites. The elevation of the marsh at site MK-4 was slightly higher than at the other three sites, and this may explain why the accretion and accumulation rates were low locally.

Overall, the agreement between ^{210}Pb and ^{137}Cs derived rates of accretion and accumulation was excellent; for a given core the rates differed by less than 10%. This suggests that sediment accumulation has been more-or-less invariant over the time frames represented by ^{210}Pb and

^{137}Cs . A noteworthy observation is the presence of measurable ^{137}Cs activity at the tops of all of the cores collected in the marsh. Given that the global atmospheric flux of ^{137}Cs has been negligible since the early 1980s, the core data implies that previously deposited ^{137}Cs has been redistributed in the system. In general, wash-in of soil-bound ^{137}Cs from the watershed and (or) erosion of ^{137}Cs -labelled subtidal deposits can account for redeposition of ^{137}Cs in river-estuaries. Additionally, bioturbation can rework ^{137}Cs from depth, but the shapes of the ^{210}Pb and ^{137}Cs profiles for all four cores are not suggestive of intense biological mixing.

Table 6 summarizes accumulation and accretion rates derived from the radionuclide measurements. We note that rates determined for the Murderkill River marsh sites are well within the range of rates previously determined for marshes fringing the tidal Delaware River and Bay estuary (Sommerfield and Velinsky, 2010); at sites spanning tidal freshwater marsh to salt marsh, the range of ^{210}Pb accretion rates is 0.3–1.3 cm/yr with a mean of 0.50 ± 0.4 cm/yr (1σ , $n=32$ cores). Similarly, the mean ^{210}Pb and ^{137}Cs accretion rate for the four sites sampled in this study was 0.60 cm/yr and 0.54 cm/yr, respectively.

In summary, cores MK-1 through MK-4 yielded reliable mass accumulation and accretion rates with good agreement between ^{137}Cs and ^{210}Pb chronological methods. Because ^{210}Pb provides a relatively longer history of sediment accumulation than ^{137}Cs , the ^{210}Pb accretion rates were used to convert sediment depth to age for the nutrient-eutrophication histories described later. Sediment-age dates corresponding to depth intervals in the cores are presented in **Tables 2-6**. These age estimates are a key output of this project as they allow us to place the nutrient data into a historical context.

C1.3: Radionuclide Inventories and Focusing Factors

Sediment inventories of excess ^{210}Pb and ^{137}Cs were computed to compare the relative amount of radionuclide deposited among coring sites MK-1 through MK-4. Reference inventories of 28 dpm/cm² for ^{210}Pb and 21 dpm/cm² for ^{137}Cs have been determined previously from direct measurements of atmospheric deposition in the U.S. mid-Atlantic region (Olsen et al. 1985; Graustein and Turekian, 1986). These reference values represent the total amount of radionuclide that could be buried at a site if supplied by atmospheric deposition alone.

To determine the extent of radionuclide focusing within the study area, focusing factors were calculated by dividing the ^{210}Pb and ^{137}Cs reference inventories by the measured inventories.

Focusing is the process by which particle-reactive substances in an environment are preferentially delivered to some depositional sites over others as a consequence of particle transport or localized scavenging processes. A focusing factor greater than unity implies that radionuclide activity has been transported or "focused" from one location to another by particle transport and deposition, or that the actual amount of radionuclide available for deposition exceeds that which can be explained by the local atmospheric flux given by the reference inventory. In the case of ^{210}Pb , tidal flooding can supply excess ^{210}Pb to a tidal marsh in addition to that derived by direct atmospheric deposition. This is because coastal and estuarine waters contain an appreciable amount of particle-bound in addition to dissolved-phase ^{210}Pb available for deposition. Conversely, in rivers, virtually all of the excess ^{210}Pb is atmospherically supplied thus focusing occurs through ^{210}Pb redistribution within the watershed. ^{137}Cs can be similarly focused, but because it is less particle reactive than ^{210}Pb it tends to desorb from particles in brackish and saline waters. Consequently, specific activities and sediment inventories of ^{137}Cs in freshwater systems tend to be higher than in estuaries and coastal waters. Hence, the physiochemical behavior of ^{137}Cs must be taken into account when interpreting ^{137}Cs focusing factors for aquatic sites that span a wide range of salinities.

Radionuclide inventories and focusing factors for the four coring sites are presented in **Table 6**. Excess ^{210}Pb inventories ranged from 47 to 66 dpm/cm², in all cases exceeding the 28 dpm/cm² reference inventory (focusing factors were 1.6–2.4). The measured inventories suggest that atmospheric deposition directly over the marsh is not the sole source of ^{210}Pb in the sediment column. ^{210}Pb inventories varied proportionally with mass accumulation rate, suggesting that the total amount of excess ^{210}Pb sequestered at the coring sites is controlled by the rate of mineral sediment accumulation as opposed to local variations in ^{210}Pb availability. In other words, the amount of tidally derived ^{210}Pb available for deposition on the marsh is similar at all of the coring sites. This is confirmed by the similar initial specific activity (A_o) of ^{210}Pb measured at the tops of the cores (~10 dpm/g).

Sediment inventories of ^{137}Cs ranged from 9.2 to 10.0 dpm/cm², less than half the reference inventory of 21 dpm/cm² (focusing factors were 0.44–0.48). These relatively low inventories most likely reflect physiochemical factors that limit ^{137}Cs uptake by particles in brackish and salt water. As was the case with ^{210}Pb , there were no along-estuary gradients in measured ^{137}Cs inventory, which implies that the sampled area is well-mixed with respect to these radionuclides.

In sum, the radionuclide inventories suggest that particle-reactive constituents entering the tidal and estuarine Murderkill River will exhibit preferential patterns of accumulate in marsh sediments that reflect factors including source location, particle affinity, and tidal transport phenomena.

C2: Nutrients/Eutrophication

C2.1: Sediment Total Carbon, Total Sediment Nitrogen and Total Sediment Phosphorus

Sediment organic carbon (SOC) concentrations for the four cores ranged from 2.3% to 34.3% on a dry weight basis (dw) with an average of $13.4 \pm 8.1\%$ SOC ($\pm 1\sigma$); **Tables 7-10; Figure 8**). Similarly, total nitrogen (TN) ranged from 0.21 to 1.7% N with an overall average of $0.68 \pm 0.37\%$; whereas total sediment phosphorus (TSP) ranged from 0.03 to 0.21% TSP with an overall average of $0.07 \pm 0.03\%$. Similar concentrations were found for TN and TSP in marsh cores taken from the river in the summer of 2007 (Chesapeake Biogeochemical Associates, 2007)

Sediment total nitrogen concentrations were highest in the bottom half of the cores from sites MK-1 and MK-2 (**Figure 8**). At MK-1, TN decreased from 50 to 40 cm and then remained fairly uniform towards the surface. A similar change occurred in MK-2 except that concentrations increased from 25 cm to the surface. In both MK-3 and MK-4, TN concentrations remained fairly constant in the entire core. A similar distribution was observed for SOC in all cores (**Figure 8**). SOC concentrations were highest in the bottom sections of the cores, MK-1 and MK-2 (up to 34%), decreasing to lower concentrations by 40 cm. In core MK-2 there was also a slight increase towards the surface above 40 cm. Concentrations of TOC in MK-3 was intermediate to those upstream and downstream (range of 3.4 to 24%). Total sediment P (TSP) increased slightly towards the surface in all cores, with the greatest increase observed in MK-4 (**Figure 8**). Concentrations of TSP were generally $< 0.1\%$ below 10 cm, increasing towards the surface.

The carbon to nitrogen ratio (C/N; atomic units) can be used as a tracer of the source of organic matter to a location and potential diagenetic changes that could occur during burial (Jasper and Gagosian, 1990; Meyers, 1994; Prahl et al., 1994). Diagenesis is any chemical, physical, or biological change undergone by sediment after its initial deposition (Bernier, 1980; Burdige, 2006). For example, terrestrial material (e.g., trees) are rich in cellulose (i.e., higher C)

compared to algae or marsh plants that have less structural material and are higher in proteins (i.e., higher N). Typical marine plants have C to N ratios of ~ 4-10 whereas terrestrial material can have C to N values > 15-20. Diagenesis of recent sediments tends to increase the C to N ratio due to preferential remineralization and release of nitrogen compounds; however, re-incorporation of bacterially derived N can increase the C to N ratio over time (Fogel et al., 1989; Benner et al.; 1991). In the upstream MK-1 core (**Figure 8**), the C/N ratio was only slightly higher at depth (18 versus ~24) suggesting preferential loss of nitrogen with burial, but at MK-2 there was no substantial change downcore (C/N of 24±2). The carbon to nitrogen ratio at MK-3 revealed slightly higher values in the surface (up to 40 at 24-26cm), decreasing with depth. Lastly, at MK-4, the C to N ratios was fairly constant with depth (18.5±2.8) except for the upper section in which the C/N was 30.4.

C2.2: Stable Isotopes of Carbon and Nitrogen

Organic carbon and nitrogen isotopic ratios are useful for distinguishing between marine and continental plant sources of sedimentary organic matter, processing and cycling of nutrients and, in some instances, the level of system-wide productivity (Fry, 2006 and others). Most photosynthetic plants incorporate carbon into organic matter using the C₃ Calvin pathway, which biochemically discriminates against ¹³C to produce a δ¹³C shift of about -20‰ to -30‰ from the isotope ratio of the inorganic carbon source. C₄ plants (e.g., corn, *Spartina*) incorporate CO₂ using a different system (PEP) that discriminates against ¹³C to produce a δ¹³C shift of about -8‰ to -15‰ from the isotope ratio of the inorganic carbon source. Organic matter produced from atmospheric CO₂ (δ¹³C ~ -7‰) by land plants and typical of a tidal freshwater wetlands, using the C₃ pathway consequently has an average δ¹³C (PDB) value of about -27‰ (O'Leary, 1988). The source of inorganic carbon for marine algae (C₃ plants) is dissolved bicarbonate, which has a δ¹³C value of about 0‰. Marine organic matter consequently has δ¹³C values between -20‰ and -22‰. The isotopic difference between organic carbon produced by C₃ land plants and marine algae has been used to trace the delivery and distribution of organic matter to sediments in estuarine and coastal areas (Cifuentes et al., 1988; Fogel et al. 1992, and many others).

Carbon isotope ratios can be affected by photosynthetic dynamics and by post-depositional diagenesis (Dean et al., 1986; Fogel et al., 1992; Canuel et al., 1995; Zimmerman and Canuel,

2002), thus isotope data must be interpreted cautiously. A big factor that can impact $\delta^{13}\text{C}$ values of plant material is the availability of CO_2 and rate of production during photosynthesis and the possibility of selective diagenesis of organic matter fractions that are isotopically heavy or light. Although the change in $\delta^{13}\text{C}$ appears to be small ($<2\text{‰}$) during recent diagenesis (Hayes et al., 1989; Meyers, 1994), shifts due to the availability and rate of production, due to nutrient enrichment or limitation and other factors, can have a large impact on the resultant $\delta^{13}\text{C}$ values of deposited organic matter (Fogel et al., 1992; Schelske and Hodell, 1995; Church et al., 2006).

Differences exist between the natural abundances of stable nitrogen isotopes ($\delta^{15}\text{N}$, $^{15}\text{N}/^{14}\text{N}$) in dissolved and particulate matter from terrestrial, estuarine, marine, and anthropogenic sources (Kendall, 1998 and others). Terrestrial-occurring soil nitrogen can have a wide range of values, but in general range from -1 to $+4\text{‰}$) similar to atmospheric nitrogen (0‰). Nitrogen isotopic compositions from marine sources tend to be slightly enriched in the heavier isotope (^{15}N) and are very dependent on the source of dissolved nitrogen and its $\delta^{15}\text{N}$, and processing in the system (e.g., ammonification, nitrification, denitrification, etc) at the time of formation. A dominant process in many aquatic environments (and groundwater) is denitrification (Cline and Kaplan, 1975). Denitrification is a microbially facilitated process of dissimilatory nitrate reduction that ultimately produces molecular nitrogen (N_2) through a series of intermediate gaseous nitrogen oxide products. This microbial process uses dissolved nitrate during oxidation of organic matter and as nitrate is consumed there is an enrichment of residual nitrate in the system. Algal/plant production and its $\delta^{15}\text{N}$ from nitrate would reflect the balance between processes and inputs. There are many points in which nitrogen can be fractionated and its isotopic composition altered. For example, wastewater from treatment facilities has been shown to increase the $\delta^{15}\text{N}$ of various fish species (Lake et al., 2001) due to the selective removal of the light isotope (^{14}N) nitrogen during treatment. Anthropogenic nitrogen was substantially enriched in watersheds with greater amount of urbanization and wastewater inputs and the nitrogen was shown to be incorporated into the aquatic food web (McCelland et al., 1997).

The isotopic compositions of sediment C and N exhibited interesting changes within each core, especially the two upper river sites (**Figure 9**). The carbon isotopic composition of the sediment ranged from -27 to -19‰ ($\delta^{13}\text{C}$ average of -17.6‰). In cores MK-1 and MK-2, the $\delta^{13}\text{C}$ was lowest in the bottom sections of the core below 40 cm ($\sim -27\text{‰}$) (**Figure 9**). In each of these two cores, the sediment organic matter become enriched in ^{13}C reaching a $\delta^{13}\text{C}$ of -17 and -

15‰ at the surface. These changes were not present in cores from MK-3 and MK-4. At both sites the $\delta^{13}\text{C}$ was fairly uniform (except in the very bottom of core MK-3), averaging $-16\pm 0.6\text{‰}$ and $-15\pm 0.6\text{‰}$, respectively.

The $\delta^{15}\text{N}$ of the sediment increased towards the surface to varying degrees in all cores (**Figure 9**). As with the carbon isotopic signatures, at sites MK-1 and MK-4 the $\delta^{15}\text{N}$ started to change at around 40 cm, increasing towards the surface. In core MK-1, the $\delta^{15}\text{N}$ was $\sim 1\text{‰}$ near the bottom increasing to 10‰ at the surface. Similarly, at MK-2, the $\delta^{15}\text{N}$ increased to approximately 8‰ near the surface. In cores MK-3 and MK-4 the increase in $\delta^{15}\text{N}$ towards the surface was less pronounced. For example, at MK-4 the $\delta^{15}\text{N}$ increased from approximately 1.8‰ in the bottom 40cm to approximately 3 to 4.5‰ at the surface.

C2.3: Diatom Analysis and Assemblages

Diatoms are microscopic, photosynthetic algae with a siliceous structure. They contain yellow-brown pigments and therefore are also referred to as golden algae. Comprising one of the most common types of algae, they are found in a diverse range of environments from freshwater to marine. Diatom species are differentiated by their shape and characteristics of their siliceous structure. The main forms are centric (i.e., circular, radial symmetry), and pennate (i.e., having bilateral symmetry). They exhibit two main living modes in the environment: planktonic and benthic (i.e., living on or in the bottom substrate).

Diatoms are one of the most powerful water quality indicators; they colonize virtually every aquatic microhabitat and many diatom species have very strict ecological requirements, with well-defined optima and tolerances for environmental variables such as pH, nutrient concentration, salinity, water transparency and physical habitat. Because of their strong relationships with environmental conditions, diatoms are used to derive inference models for such environmental factors. The inference models are developed using calibration sets of both diatoms and measured environmental variables for specific geographic regions. To produce robust quantitative models, the calibration sets require at least 30 sampling sites that maximize the gradient length covered by the variable of interest (e.g., phosphorus concentration, pH, etc.). These models can then be used to infer the environmental parameter of interest when instrumental measurements are not available, and have been successfully used to reconstruct

reference conditions and assess the impact of anthropogenic activities on aquatic systems (e.g., Cooper et al., 1993; 1995; Bennion et al., 2001).

Diatom assemblages have been shown to be important indicators of nutrient concentration within freshwater and marine environments. Due to the fact the diatoms respond quickly and directly to nutrients, they have been used for many years as indicators of nutrient changes in aquatic systems (Potapova et al., 2004; Potapova and Charles, 2007; Ponader et al., 2008). In this study we take a first approach using diatom metrics calculated using species-ecological models developed by van Dam et al. (1994). While this data set is mostly based on freshwater species it does contain some marine diatoms. It provides ecological information for both diatoms and water characteristics is not currently available for the Delaware Estuary.

A total of 56 sections were analyzed for diatom composition from four Murderkill River cores (MK-1, MK-2, MK-3, and MK-4). At least 400 valves were counted for each sample and over 300 taxa (**Appendix I**) were identified in these samples allowing a robust analysis for nutrient and ecological conditions. The distribution of key diatom species (those occurring with more than 5% relative abundance) in each of the cores is presented in **Figures 10-13**. Brief descriptions of the species' stratigraphic distribution and major shifts in diatom assemblages are provided below for each core.

Core MK-1: 215 species were identified in 15 sections from core MK-1, of which only 19 species occurred with abundances > 5% in at least one sample. The most abundant species were: *Stauroforma exiguiformis* (maximum relative abundance 66%), *Cymatosira belgica* (32%), *Navicula salinarum* (17%), *Denticula subtilis* (16%), *Nitzschia brevissima* and *N. frustulum* (10%). In this core two distinct types of diatom assemblages were present: one with the species *Stauroforma exiguiformis* dominant and rarer benthics such as *Nitzschia nana*, *Pinnularia subcapitata* occurring between the core bottom and 42-44 cm core depth. This freshwater, circumneutral to slightly acidic type of diatom assemblage was replaced above 42-44 cm by an assemblage with a much more diverse species composition, mostly represented by brackish-marine taxa such as *Cymatosira belgica*, *Navicula salinarum*, *Denticula subtilis*, *Nitzschia brevissima*, *Cyclotella striata* (**Figure 10**). The major shift to brackish-marine diatom species took place in the mid to late 1950s.

Core MK-2: 172 species were identified in 14 sections from this core with 21 species > 5% relative abundance in at least one sample. The most abundant species were: *Stauroforma exiguiformis* (maximum relative abundance 71%), *Pseudostaurosira subsalina* (28%), *Navicula salinarum* (22%), and *Diploneis smithii* (22%) (**Figure 11**). Similar to core MK-1, a major shift from freshwater-type diatoms with abundant *S. exiguiformis* to marine-brackish species took place at 36 cm core depth. The freshwater-type of assemblage differed slightly from core MK-1 by the presence of abundant *D. smithii* amongst other benthic species such as *Pinnularia viridis* and *Calloneis bacillum*) at the core bottom, and species that are usually present in more acidic waters such as *Frustulia saxonica*, and *Eunotia* spp. The major change in diatom assemblages from freshwater to marine took place in the late 1940 to mid 1950s.

Core MK-3: In this core, 152 species were identified, of which 21 species were present with relative abundances > 5%. The most abundant species were: *Luticola mutica* (maximum relative abundance 43%), *Caloneis bacillum*, *Paralia sulcata* (14%), *Denticula subtilis* (13%), *Navicula cincta* (12%), *Actynocyclus normanii* fo. *subsalsum*, and *Navicula* sp. 1 MUR DME (10%) (**Figure 12**). Below 28 cm depth of the core, diatom species were very diverse with nutrient-rich freshwater species (*Calloneis bacillum*, *Nitzschia frustulum*) as well as brackish-marine species (*Denticula subtilis*, *Paralia sulcata*). Above 28 cm, a major shift took place with *Luticola mutica*, a subaerial diatom species usually found on rocks and walls, becoming abundant and even dominant in many core sections of this interval. In the upper two sections (0-2 and 4-6-cm intervals), *L. mutica* decreased while marine species such as *A. normanii*, *Cyclotella striata*, and *Cymatosira belgica* increased in relative abundance.

Core MK-4: A total of 136 species were identified in 11 sections from this core. Diatom assemblages were very diverse, composed of both marine-brackish and freshwater eutrophic species. No species were dominant in this core except for *Cymatosira belgica* in the top core interval (39%). No particular trend was identified in this core's diatom assemblages either, except that the coastal marine *Cymatosira belgica* displayed increased abundances in both the bottom and upper intervals (**Figure 13**).

The samples collected in the Murderkill River wetlands span a range of marine-brackish to freshwater diatom species, making detailed analysis complex. To extract the ecological information contained in the diatom data identified in this region, various metrics were calculated for salinity, pH, nitrogen uptake metabolism, and the trophic state (van Dam et al., 1994; USGS NAWQA; Hall and Smol, 1999). These metrics are often used to assess freshwater systems rivers and streams in the U.S. All of these metrics have strengths and weaknesses that vary with the quality of the ecological information available for the taxa found in each of the cores. Because most marine or brackish species present in the Murderkill River region do not occur in van Dam et al. (1994) data sets, metric calculations were based mostly upon the freshwater species occurring in these cores. The van Dam et al. (1994) summary and synthesis of ecological characteristics of taxa is based primarily on samples from Western Europe, and generally includes taxa with wide geographic distributions. The van Dam et al. (1994) autecological values for many taxa have been revised to be more relevant for the U.S., and are contained in a data compilation developed for the USGS National Water Quality Assessment (NAWQA) program by Porter (2008) and Porter et al. (2008). To allow a comparison between the current study in the tidal Murderkill River and the previous studies of the tidal Christina River and Saint Jones River (Velinsky et al., 2007; 2010), the data set and metrics by van Dam et al. (1994) will be the focus of this discussion.

The van Dam et al. (1994) diatom metrics show an abrupt increase in the proportion of eutrophic species (i.e., high nutrient species) in cores MK-1, MK-2, and MK-3, starting at 42-44 cm, 34-36 cm, and 18-20 cm towards the surface of the core, respectively (**Figure 14; Tables 11-14**). Below this depth range, oligotrophic, freshwater species are present in MK-1 and MK-2, while in MK-3, diatom species are mostly mixed meso- and eutrophic species, with a small spike in oligotrophic species at 70 cm depth. Core MK-4 mainly displays an increasing (but fluctuating) trend in eutrophic species, except for the top two intervals.

To evaluate how nutrient conditions, as reflected by total sediment N and P, impacted the diatom species and metrics in each core, N and P were plotted against the metrics for eutrophic species (**Figures 15-16**). For cores MK-1 and MK-2, there was an inverse relationship ($r^2=0.74$ and $r^2=0.73$ for MK-1, and MK-2, respectively) between eutrophic species (as per Van Dam et al., 1994) and TN (i.e., the higher the TN, the lower the %eutrophic diatoms; **Figure 15**). In core MK-3 there was a slight positive trend while there was no relationship observed in MK-

4. In all cores, there was no relationship between TSP and the diatom trophic metric (**Figure 15**). The accumulation rates of N and P (see below) were calculated and plotted against the %eutrophic diatoms as this might be a better integrator of N and P inputs to the river. As with the concentration data there was no consistent relationship between the N or P accumulation rates (over time) and changes in the trophic status of the tidal river.

The trends observed in diatom autecology and sediment stable isotopes reveal the complexity of nutrient inputs and biogeochemical processes taking place within the tidal Murderkill River, and suggest that more elaborate analyses of data are necessary to reveal their dynamics. For example, cross-correlation analyses are necessary to account for the lag response between changes in nutrients and shifts in diatom species, or to account for lags related to sedimentary nutrient re-mobilization. However, within the Murderkill river system, the upper part of cores MK-1 and MK-2 reveal a shift in diatom species related to increasing nutrient concentration which clearly correlates with the positive trend observed in $\delta^{15}\text{N}$. These changes start to occur in the late 1940s and may reflect increasing agricultural inputs and population growth in the watershed. A similar, but much more recent change was observed in core MK-3, while core MK-4 did not reveal a particular trend in nutrient concentration.

C3: Historical Analysis

A major focus of this study was to determine the changes in the loadings of nitrogen and phosphorus over time within the tidal Murderkill River system. These changes are thought to have occurred in the early to mid 1970s with the introduction of wastewater from the KC WWTP that came online during this time period. In addition, an objective was to determine if there was any indicator of an ecosystem change (e.g., change in plant flora, oxygen dynamics, marsh production, etc.) as a result of the facility and effluent discharges of nitrogen and phosphorus. Lastly, we wanted to determine if the marshes along the tidal river are a sink for N and P via sediment burial and how this relates to other inputs.

It is important to recognize that both nitrogen and phosphorus can undergo substantial biogeochemical processing and diagenetic changes during burial (**Figure 1**). In brief, inorganic forms of N and P are taken up by marsh plants, phytoplankton and benthic algae and incorporated into marsh sediments. During burial, microbial activity (both oxic and anoxic) can release organic nitrogen and phosphorus into the porewaters of the sediments which can move

back into the overlying waters. A major sink or loss reaction for nitrogen is denitrification which converts oxidized nitrogen to nitrogen gas which is then removed from the system. These nutrients, along with externally introduced nutrients, are then transported downstream or taken up again during photosynthesis. This recycling of nutrients in a marsh is a major process that impacts overall marsh/estuarine productivity and transport to coastal areas. In this regard, depending on the environment and specific characteristics, such as the magnitude of nutrient loadings, sedimentation rates, oxic/anoxic conditions, it may or may not be possible to measure changes in nitrogen or phosphorus accumulation rates that reflect inputs from external sources such as with chemicals that do not undergo significant biogeochemical reactions (e.g., PCBs, Pb, Zn, etc).

C3.1. Changes in Nutrients and Diatoms in Murderkill River Cores

In the tidal Murderkill, total nitrogen in sediments over time varied among core locations (**Figure 17**). At the upper coring site (MK-1), TN concentrations were highest between 1900 and 1920 (~ 1.5% N), decreasing to approximately 0.6% N by 1940 and remaining constant to the present. A similar change occurred at site MK-2, but the concentrations started to increase after approximately 1970, around the time the KC WWTP came online. Site MK-2 was on the marsh directly across the river from the tidal creek into which the treatment plant discharges. At site MK-3 and farther down-estuary at site MK-4, TN concentrations were fairly constant from 1900 to the present. However, concentrations in the near surface sections of cores were highest at site MK-2 and decreased down-estuary.

The stable isotopes of sediment nitrogen may give some indication as to changes in the source of nitrogen and biogeochemical cycling within the tidal river. Ulseth and Hershey (2005), Leavitt et al. (2006) and Velinsky (unpublished data) show that higher $\delta^{15}\text{N}$ of the nitrogen is associated with higher inputs from urban sources and waste water. Bratton et al. (2003) showed similar trends in cores from the Chesapeake Bay. Lastly, in Woodbury Creek and Oldmans Creek marshes in New Jersey, an increase in $\delta^{15}\text{N}$ observed in sediment cores was attributed to both increased nitrogen loading and changes in the way nitrogen is processed in wastewater treatment plants (Church et al., 2006). Downcore variations of $\delta^{15}\text{N}$ are somewhat difficult to interpret given the number of biogeochemical processes influencing nitrogen. In the Murderkill River marshes, there was a pronounced increase in the $\delta^{15}\text{N}$ of the sediment nitrogen in cores

MK-1, MK-2 and MK-3 and to a lesser degree in MK-4 (**Figure 17**). The increase in the $\delta^{15}\text{N}$, from $< 1 \text{ ‰}$ to between 3 and 10 ‰ at the surface, started in the 1950s. This occurred long before the changes in nitrogen loadings from the addition of the KC WWTP (mid-1970s). The low $\delta^{15}\text{N}$ prior to the 1950s, especially in the upper estuary cores (MK-1, 2 and 3), suggest more terrestrial sources (Kendall, 1998) of N, which would be around 0 to +2‰, whereas the higher $\delta^{15}\text{N}$ values downstream at MK-4 are influenced by marine sources which tend to have higher $\delta^{15}\text{N}$. Similarly, Elliot and Brush (2006) showed a correlation between the $\delta^{15}\text{N}$ of tidal wetland core sediments and estimates of nitrogen wastewater loadings over time. However, it is unclear what may have occurred in the Murderkill watershed pre-1970s to substantially increase the $\delta^{15}\text{N}$, but it is most likely a combination of changes such as increasing population growth and septic system discharges, agricultural practices and changes in nitrogen fertilizers. Interestingly, surface sediment $\delta^{15}\text{N}$ decrease from 10 ‰ at the upper station (MK-1) and decrease to MK-4, again reflecting the changes in nitrogen sources from more land derived to more marine, Delaware Bay, derived sources.

Concentrations of total sediment phosphorus (TSP) changed over time at all coring locations to some degree (**Figure 18**), especially at site MK-4. At MK-1, concentrations averaged $0.073 \pm 0.007 \text{ ‰P}$ between 1900 and 2003, then slightly increasing at the surface to 0.11 ‰P. There was a slight change at the same time as the shift in $\delta^{13}\text{C}$ of the sediment. At site MK-2 concentrations were similar until approximately 1980 at which time concentrations increased 0.12‰P. A similar trend, but at lower concentrations, was observed at MK-3 with concentrations reaching only 0.09‰P at the surface. The largest change was observed further down-estuary at MK-4 (**Figure 18**). At this location, TSP was constant from the 1900s to 1970 after which concentrations substantially increased to 0.15 to 0.20‰P in 2008. Although the increase in concentrations at sites MK-2 and MK-3 may be a result of the discharge starting in the 1970s, it is unclear why the largest increase was observed much farther down-estuary at MK-4. It is possible that drainage of water from the large wetlands area upstream of Old Brockonbridge Gut (near South Bowers Beach; **Figure 1**) imparts a large load of phosphorus to the main river and that a portion of this phosphorus is deposited on the marsh platform adjacent to the tidal gut.

Schelske and Hodell (1991; 1995), and Perga and Gerdeaux (2004) have shown a relationship between the concentration of P in the sediments or water column and the isotopic composition of carbon ($\delta^{13}\text{C-OC}$) in the sediments or in fish scales over time, respectively. The

fish scales were an integrator of the base of the food web (i.e., phytoplankton) and reflect nutrient inputs. Similarly, Church et al. (2006) showed a relationship between sedimentary P and $\delta^{13}\text{C-OC}$ in a marsh core from Woodbury Creek, New Jersey. It is possible that as P levels increase, primary productivity increases to a point at which there is reduced isotopic fractionation during enzymatic uptake of dissolved CO_2 . This reduced fractionation would result in higher isotopic compositions of organic matter (i.e., more ^{13}C -enriched) and would suggest that P was helping to control aquatic productivity. Therefore, if phosphorus is limiting production, the $\delta^{13}\text{C}$ of the organic carbon in a core may reflect a system-wide change in productivity. In the Murderkill River, at only the upper two locations (MK-1 and MK-2) was there a significant enrichment of ^{13}C in core sediments over time. Sediment $\delta^{13}\text{C}$ changed from approximately -25‰ to \sim -15‰ from the 1900s to the present. In both downstream cores there was only a slight change throughout the entire core with an overall average of -15.4 ± 1.1 ‰. There was no relationship between TSP and $\delta^{13}\text{C}$ of the organic carbon, suggesting that phosphorus may not be limiting in this system and that other factors are controlling the $\delta^{13}\text{C}$ of the organic carbon in the sediments (see below). It is possible that changes in the source of organic matter to the marshes (i.e., terrestrial-upland material versus *in situ*-local production), or that *in situ* marsh processes and plant communities may have changed over time.

The diatom metrics used to assess eutrophication for each core are plotted against time in **Figure 19**. It appears that there are two separate trends occurring in the cores from the various sites over time. At sites MK-1 and MK-2 there is a distinct increase in the index for eutrophic conditions starting in the late 1930s to early 1940s. The index then decreased in the late 1950s and increased slightly after the 1970s. At MK-3, there was no increase in the late 1930s, but there was a gradual increase in the late 1960s to early 1970s which maybe an indicator of the increase in nitrogen and phosphorus loadings from various sources, including the KC WWTP. There was no substantial change in the eutrophic index at MK-4 over time.

One major change that occurred in cores from sites MK-1 and MK-2 was a shift from freshwater species of diatoms to more marine/estuarine species starting in the 1950s (**Figure 10 and 11**). That is, a major shift from freshwater-type diatoms (e.g., *S. exiguiformis*) to marine-brackish species. Other freshwater-benthic diatoms included abundant *D. smithii*, and *Pinnularia viridis* and *Calloneis bacillum* at the core bottom. Species common in more acidic waters such as *Frustulia saxonica* and *Eunotia* spp were also present. These were replaced by

brackish-marine taxa such as *Cymatosira belgica*, *Navicula salinarum*, *Denticula subtilis*, *Nitzschia brevissima*, and *Cyclotella striata*. This shift was not evident farther downstream at sites MK-3 and MK-4.

C3.2 Changes in Marsh Ecology

In addition to the change in diatom flora, there was a large shift in the stable isotopic composition of carbon and nitrogen during this time period (i.e., 1930s to the present; **Figures 17 and 18**). The changes in the $\delta^{13}\text{C}$ composition of sedimentary organic carbon over time indicate that the marsh shifted from C3 dominant plants (e.g., wild rice, spatterdock, and others) to C4 dominant plants (e.g., *Spartina* sp.). *Spartina* species have $\delta^{13}\text{C}$ values enriched in ^{13}C , characteristic of plants using the C4 photosynthetic pathway (Smith and Epstein, 1971). Separate analysis of *Spartina alterniflora* from the Murderkill River yielded a $\delta^{13}\text{C}$ of $-13.7 \pm 0.4\text{‰}$ (n=4), close to the values exhibited in the surface sediment of all cores. Although there are some C3 saltmarsh plants (e.g., *Scirpus* sp.) that have been shown to have more negative $\delta^{13}\text{C}$ values (Cloern et al., 2002; Chmura and Aharon, 1995), it is unclear how dominant this plant was in the past. Plants typically present in a tidal freshwater marsh (e.g., wild rice, *Typha*, spatterdock, and others) use the C3 photosynthetic pathway and have $\delta^{13}\text{C}$ values enriched in ^{12}C (i.e., more negative $\delta^{13}\text{C}$) (Cloern et al., 2002; Chmura and Aharon, 1995). Presence of freshwater diatoms and the more negative $\delta^{13}\text{C}$ composition of the sedimentary organic matter in the bottom section of these two cores suggest that before 1940, these two marshes were situated in predominately tidal freshwater. Byrne et al. (2001) showed that changes in water storage and diversion in the San Francisco Estuary resulted in changes in diatoms and carbon isotopic composition of a sediment core from a tidal marsh. A similar change could be occurring in the Murderkill River.

The exact cause of this shift is unclear, but it is hypothesized that the hydrology changed in the tidal river such that more saline water moved upstream over time. Presently, salinities range from 1-5 psu to 10-15 psu over the year at the Route 1 Bridge, just upstream from MK-1 (USGS monitoring data, unpublished). It is possible that the salinities in the upper region were fresher in the past (see for example Byrne et al., 2001). The increase in salinity around the two upstream sites initiated a shift from a marsh dominated by C3 plants (i.e., more typical of a tidal freshwater wetland) to a salt marsh with a dominance of *Spartina*, that took place from the 1940s to 1960s.

There are at least two possibilities that may have caused this change: damming of upstream flows and dredging in the tidal river. In addition, there was substantial mosquito ditching during the early to mid-1930s in the lower part of the tidal river (see Bourn and Cottam, 1950; Clarke et al., 1984; LeMay, 2007 for impacts). While there is little information concerning the timing and extent of tributary damming upstream, there are three dams: McGinnis Pond on Hudson Branch, Killen and Coursey Pond on Upper Murderkill River, and Andrews Lake on Pratt Branch. It is thought that these ponds are old mill ponds created to produce local energy and were created prior to the 1900; however, more information is needed to confirm the timeframe. With regards to dredging there is only preliminary information that dredging occurred from the mouth of the river with Delaware Bay up to Fredrica, a distance of ~11 km. Dredging the river would have allowed more saline water to move farther upstream. In either case, the marshes upstream at MK-1 and MK-2 experienced a change from a freshwater biota to a more marine biota, and this change needs to be investigated further.

C3.3: Nitrogen and Phosphorus Accumulation Rates

Sediment cores can be used to calculate the mass of nitrogen (N) and phosphorus (P) accumulating in the marsh over time. Using the mass accumulation rate and concentration of nitrogen and phosphorus (% TN and %TSP) in each interval, the N and P accumulation rate (mg N or P/cm²-yr) for each year can be obtained. It should be noted that missing intervals that were not analyzed for N and P were linearly interpolated from the interval concentrations above and below. The accumulation rates can then be applied to estimate of wetland area to provide an estimate of the amount of N and P removed (via burial) by wetland processes. Other removal/transport processes (e.g., denitrification, recycling, sediment movement) would need to be determined to provide a complete N and P cycle in the river.

Nitrogen accumulation rates for the four locations presently range from approximately 1 to 2.5 mg N/cm²-yr (**Figure 20**). At sites MK-1, MK-2 and MK-3, rates were ~ 2 mg N/cm²-yr before the 1960s decreasing to a minimum in 1970 then increasing from approximately 1970–1975 to the present. At site MK-4 there was no minimum but a small increase over time to the present. Phosphorus accumulation rates exhibited a similar pattern with time (**Figure 20**), except that there was no accumulation minimum present (except possibly MK-2). In general, starting after approximately 1975, P accumulation increased from 0.05 to 0.1 mg P/cm²-yr to

0.15 to 0.3 mg P/cm²-yr at the present. These rates and the change over time are similar to those found in a tidal freshwater wetland in Patuxent River (Kahn and Brush, 1994) and in marshes along the Patuxent salinity gradient (Hartzell et al., 2010).

A rough estimation of the area of tidal wetlands fringing Murderkill River is 4,300 acres (1.74 X 10⁷ m²; DNREC, 2005). Using this area and the core-top values for N and P accumulation yields current burial rates (gross rates) of 25 X 10⁶ g N/month and 3.4 X 10⁶ g P/month, respectively. Ullman et al. (2010) estimated monthly N and P external loading to the tidal Murderkill River from the KC WWTP, upstream loads from the watershed and direct atmospheric loads. Nitrogen loadings (predominately from the upper watershed) ranged from approximately 20 to 100 X 10⁶ g N/month while P loads (predominately from the KC WWTP) ranged from < 0.5 to 13 X 10⁶ g P/month. These calculations show that marsh accumulation can sequester a majority of the P and N loads from the various sources (i.e., point sources for P and non-point sources for N). However, sediment recycling of N and P (Bernier, 1980; Burdige, 2006) are not accounted for in these estimates and will modify and most likely reduce these fluxes. In this regard, Ullman et al. (2010) estimated that the tidal river may not be attenuating phosphorus substantially, whereas nitrogen may be significantly removed through denitrification and burial within the estuary and not exported to Delaware Bay. Nonetheless, the estimates provided above show that the marshes have a potential to trap both N and P before being exported to the tidal river.

D) Summary and Conclusions

This study involved the chemical analyses of sediments subsampled from four sediment cores taken from the tidal and estuarine region of the Murderkill River in Kent County, Delaware. The objective was to determine the chronology of carbon, nitrogen, and phosphorus deposition, nutrient loading histories, and related ecological responses based on preserved diatom assemblages and stable isotope signatures of carbon ($\delta^{13}\text{C-OM}$) and nitrogen, ($\delta^{15}\text{N-TN}$). Cores were collected from marsh sites upstream near Frederica, to downstream near Bowers Beach, and dated using ^{210}Pb and ^{137}Cs radiometric methods. All cores provided sufficient temporal coverage (> 50 yr) for detailed chemical analyses.

Major findings of this study include:

- Sediment mass accumulation rates, determined by ^{210}Pb and ^{137}Cs geochronology, ranged from 0.10 to 0.20 $\text{g}/\text{cm}^2/\text{yr}$, and corresponding accretion rates ranged from 0.31 to 0.74 cm/yr . These rates are comparable to those measured in other tidal marshes of the Delaware Estuary.
- Sediment concentrations of C, N and P are typical for tidal fresh and marine marshes throughout the Delaware Estuary. Sediment organic carbon (SOC) concentrations ranged from 2.3% and 34.3% dw, total nitrogen (TN) ranged from 0.21 to 1.7% N and total sediment phosphorus (TSP) ranged from 0.03 to 0.21% P. Sediment nitrogen concentrations increased in only one site (MK-2), while TSP concentrations showed only a small increase towards the surface at three sites, but a larger increase in MK-4, the most downstream location. Sediment organic carbon exhibited a substantial decrease towards the surface at MK-1 and 2 in the upper river.
- The analyses of the diatom assemblages and metrics indicate a shift toward more eutrophic species starting in the late 1940-1950s (cores MK-1 and MK-2) and 1970s (core MK-3), but no clear trend was observed in core MK-4. There was no significant relationship between the diatom metrics and levels of N and P.
- In cores MK-1 and MK-2, there was a diatom flora change from freshwater to brackish-marine species with time, along with a change in the stable isotopic composition of sedimentary organic carbon. The organic carbon shifted from more negative values to

more positive values indicating a change from C3 plants to C4 plants with time starting in the late 1930s to early 1940s. This suggests that the average salinity of water in the upper tidal river has increased in recent times.

- Sediment N and P accumulation rates increased two fold since 1975-1980. This is approximately the same time frame as the onset of effluent discharge from the KC WWTP.
- Preliminary estimates suggest that current burial rates in the marsh can remove a substantial fraction of both N and P; however, recycling rates need to be considered for more accurate estimates of net burial rates. This study provides important data and results that can be incorporated into an ecosystem model for the tidal river to help determine potential water quality changes in the future.

E) Acknowledgments

We would like to thank Mihaela Enache, Don Charles, Paul Kiry, Dack G. Stuart and Paula Zelanko for field and laboratory assistance and data interpretation. Dr. Mihalea Enache did diatom counts, calculated diatom metrics and provided assistance with report preparation. We appreciate the help of Roger Thomas and Jared Halonen in field. Robin Davis provided assistance with final report preparation and review.

F) References

- American Public Health Association, American Water Works Association and Water Environment Federation (APHA, AWWA and WEF). 1995. *Standard Method for the Examination of Water and Wastewater*, 19th Edition. Washington, DC.
- Aspila, K.I, Aspila, H., Agemian A. and Chau, S.Y.,1976. A semi-automated method for the determination of inorganic and total phosphate in sediments. *Analyst* 101: 187-197.
- Aurand, D. and F.C. Daiber. 1973. Nitrate and nitrite in the surface waters of two Delaware salt marshes. *Chesapeake Science* 14: 105-111.
- Bennion, H., P.G. Appleby, and G.L. Phillips. 2001. Reconstructing nutrient histories in the Norfolk Broads: implications for the application of diatom-phosphorus transfer functions to shallow lake management. *Journal of Paleolimnology* 26:181-204.
- Berner, R. 1980. *Early Diagenesis: A Theoretical Approach*. Princeton University Press, 241 pp.
- Benner, R., M. L. Fogel, and E. K. Sprague. 1991. Diagenesis of belowground biomass of *Spartina alterniflora* in salt-marsh sediments. *Limnol. Oceanogr.* 36: 1358-1374.
- Bratton, J.F., S.M. Coleman and R.R. Seal, II. 2003. Eutrophication and carbon sources in Chesapeake Bay over the last 2700 yr.: Human impacts in context. *Geochim. Cosmochim. Acta.* 67: 3385-3402.
- Brush, G.S. and L.M. Brush. 1994. Transport and deposition of pollen in an estuary: signature of the landscape. In: edited by A. Traverse, *Sedimentation of Organic Particles*, Cambridge University Press; Great Britain.
- Brush, G.S., E.A. Martin, R.S. DeFries, and C.A. Rice 1982. Comparisons of ^{210}Pb and pollen methods for determining rates of estuarine sediment accumulation. *Quaternary Research.* 18: 196-217.
- Burdige, D.J. 2006. *Geochemistry of Marine Sediments*. Princeton University Press. 630 pp.
- Bourn, W.S. and C. Cottam. 1950. Some biological effects of ditching tidewater marshes. US Department of Interior, Fish and Wildlife Ser. Res. Rep. 19.
- Boynton, W.R., J.H. Garber, R. Summers, and W.M. Kemp. 1995. Inputs, transformations and transport of nitrogen and phosphorus in Chesapeake Bay and selected tributaries. *Estuaries* 18: 285-314.
- Byrne, R., L. Ingrama, S. Starratta, F. Malamud-Roama, J.N. Collins and M. E. Conrad. 2001. Carbon-isotope, diatom, and pollen evidence for Late Holocene salinity change in a brackish marsh in the San Francisco Estuary. *Quart. Res.* 55: 66-76.

- Canuel, E.A., J.E. Cloern, D.B. Ringelberg, J.B. Guckert and G.H. Rau. 1995. Molecular and isotopic tracers used to examine sources of organic matter and its incorporation into the food webs of San Francisco Bay. *Limnol. Oceanogr.* 40: 67-81
- Castro, M.S., C.T. Driscoll, T.E. Jordan, W.G. Reay and W.R. Boynton. 2003. Sources of nitrogen to estuaries of the United States. *Estuaries and Coasts* 26: 803-814.
- Charles, D. F., Knowles, C., and Davis, R. S. (Eds.). 2002. Protocols for the Analysis of Algal Samples Collected as Part of the U.S. Geological Survey National Water-Quality Assessment Program. Patrick Center for Environmental Research-Phycology Section, The Academy of Natural Sciences of Philadelphia. Report No. 02-06. 124 pp.
- Chesapeake Biogeochemical Associates. 2007. Nutrient Flux Study – July 2007 Results from the Murderkill River-Marsh Ecosystems. Initial Report submitted to Kent County Levy Court (Nov. 2007).
- Church, T.M., C. Sommerfield, D.J. Velinsky, D. Point, C. Benoit, D. Amouroux, D. Plaa and O. Donard. 2006. Marsh sediments as records of sedimentation, eutrophication and metal pollution in the urban Delaware Estuary. *Marine Chemistry* 102(1-2):72-95.
- Chmura, G.L. and P. Aharon. 1995. Stable carbon isotope signatures of sedimentary carbon in coastal wetlands as indicators of salinity regime. *J. Coast. Res.* 11:124-135.
- Cifuentes, L.A., J.H. Sharp and M.L. Fogel. 1988. Stable carbon and nitrogen isotope biogeochemistry of the Delaware estuary. *Limnol. Oceanogr.* 33: 1102-1115.
- Clarke, J.A., B.A. Harrington, T. Hruby, and F.E. Wasserman. 1984. The effect of ditching for mosquito control on salt marsh use by birds in Rowley, Massachusetts. *J. Field Ornithol.* 55: 160-180.
- Cline, J.D. and I.R. Kaplan. 1975. Isotopic fractionation of dissolved nitrate during denitrification in the eastern tropical north pacific ocean. *Marine Chemistry* 3: 271-299.
- Cloern, J.E. E.A Canuel and D.Harris. 2002. Stable carbon and nitrogen isotope composition of aquatic and terrestrial plants of the San Francisco Bay estuarine system. *Limnol. Oceanogr.* 47: 713-729.
- Cooper, S.R. 1999. Estuarine paleoenvironmental reconstructions using diatoms. in Stoermer, E.F. and J.P. Smol, editors. *The diatoms: Applications for the environmental and earth sciences*. Cambridge University Press, Cambridge, pages 352 - 373
- Cooper, S.R. 1995. Chesapeake Bay watershed historical land use: impact on water quality and diatom communities. *Ecological Applications* 5: 703-723.
- Cooper, S. R. and G.S. Brush. 1993. A 2,500-year history of anoxia and eutrophication in Chesapeake Bay *Estuaries* 16: 617-626.

Conley, D.J. and C.L. Schelske. 2001. Biogenic Silica. In: Smol, J.P., H. J.B. Birks and W.M. Last. *Tracking Environmental Change using Lake Sediments. Volume 3: Terrestrial, Algal and Siliceous Indicators*. Kluwer Academic Publishers, The Netherlands, pages: 281-294.

Cornwell, J.C., D.J. Conley, M. Owens, and J.C. Stevenson. 1996. A sediment chronology of eutrophication of Chesapeake Bay. *Estuaries* 19:488-499.

Cutshall, N.H., I.L. Larsen and C.R. Olsen. 1983. Direct analysis of ^{210}Pb in sediment samples: Self-absorption corrections. *Nuclear Instruments and Methods in Physics Research*, 206: 309-312.

Dean, W.E., Arthur, M.A., and Claypool, G.E., 1986. Depletion of ^{13}C in Cretaceous marine organic matter: source, diagenetic, or environmental signal? *Mar. Geol.*, 70:119-157.

De Michele, E. 1972. The effects of conservative and non-conservative substances on the Murderkill River. Final Rep., Dept. Nat. Res. Environ. Control; State of De. 101p.

deWitt, P. and F.C. Daiber. 1974. The hydrography of the Murderkill Estuary, Delaware. *Ches. Sci.* 15: 84-85.

DNREC. 2007. State of Delaware 2006 Combined Watershed Assessment Report (305b) and Determination for Clean Water Act Section 303(d) List of Waters needing TMDLs. State of Delaware Department of Natural Resources and Environmental Control Division of Water Resources, Watershed Assessment Section. Dover, DE.

DNREC 2006. Fact Sheet (November 1, 2006). Surface Water Discharge Section; Dover, DE.

DNREC, 2005. Technical Analysis for Amendment of the 2001 Murderkill River TMDLs. State of Delaware Department of Natural Resources and Environmental Control Division of Water Resources, Watershed Assessment Section. Dover, DE (Updated March 1, 2005).

Elliott, E.M. and G.S. Brush. 2006. Sedimented organic nitrogen isotopes in freshwater wetlands record long-term changes in watershed nitrogen source and land use. *Environ. Sci. Technol.* 40: 2910–2916.

Fogel, M.L., L.A. Cifuentes, D.J. Velinsky and J.H. Sharp. 1992. The relationship of carbon availability in estuarine phytoplankton to isotopic composition. *Marine Ecol. Prog. Ser.* 82: 291-300.

Fogel, M.L., E.K. Sprague, A.P. Gize and R.W. Frey. 1989. Diagenesis of organic matter in Georgia salt marshes. *Estuarine, Coastal Shelf Sci.*, 28: 211-230.

Follmi, K.B. 1996. The phosphorus cycle, phosphogenesis and marine phosphate-rich deposits. *Earth Science Reviews* 40: 55-124.

- Froelich, P.N., M.L. Bender, N.A. Luedtke, G.R. Heath, and T. De Vries. 1982. The marine phosphorus cycle. *Amer. Jour. Sci.* 282: 474-511.
- Fry, B. 2006. *Stable Isotope Ecology*. Springer Science+Business Media, LLC, New York.
- Graustein, W.C. and K.K. Turekian. 1986. ^{210}Pb and ^{137}Cs in air and soils measure the rate and vertical distribution of aerosol scavenging. *J. Geophys. Res.*, 91: 14,355-14,366.
- Hall, R. I., and J. P. Smol. 1999. Diatoms as indicators of lake eutrophication. Pages 128-168 in: E. F. Stoermer, and J. P. Smol (editors). *The diatoms: applications for the environmental and earth sciences*. Cambridge University Press, Cambridge, UK.
- Hartzell, J.L., T.E. Jordan, and J.C. Cornwell. 2010. Phosphorus burial in sediments along a salinity gradient of the Patuxent River, a sub-estuary of the Chesapeake Bay. *Estuaries and Coasts* 33: 92-106.
- Hayes, J.M, Popp, B.N., Takigiku, R., and Johnson, M.W., 1989. An isotopic study of biogeochemical relationships between carbonates and organic carbon in the Greenhorn Formation. *Geochim. Cosmochim. Acta*, 53:2961-2972.
- Heiri, O., A.F. Lotter and G.Lemcke. 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *Journal of Paleolimnology*. 25: 101-110.
- Hornberger, M.L., S.N. Luoma, A.Green, C. Fuller, and R. Anima. 1999. Historical trends of metals in the sediments of San Francisco Bay, California. *Marine Chemistry*. 64:39-55.
- Inn and others 2001. The NIST natural-matrix radionuclide standard reference material program for ocean studies. *Journ. Radioanal. Nucl. Chem.* 248: 227-231.
- Jasper, J.P. and R.B. Gagosian. 1990. The sources and deposition of organic matter in the Late Quaternary Pigmy Basin, Gulf of Mexico. *Geochim. Cosmochim. Acta*. 54: 1117-1132.
- Jordan, T.E., J. C. Cornwell, W.R. Boynton, and J.T. Anderson. 2008. Changes in phosphorus biogeochemistry along an estuarine salinity gradient: The iron conveyer belt. *Limnol. Oceanogr.* 53: 172-84.
- Kahn, H. and G.S. Brush. 1994. Nutrient and metal accumulation in a freshwater tidal marsh. *Estuaries* 17: 345-360.
- Kendall, C. 1998. Tracing nitrogen sources and cycling in catchments. In: *Isotope Tracers in Catchment Hydrology*; Elsevier, Amsterdam, pp. 519-576.
- Lake, J.L., R.A. McKinney, F.A. Osterman, R.J. Pruell, J. Kiddon, S.A. Ryba and A.D. Libby. 2001. Stable nitrogen isotopes as indicators of anthropogenic activities in small freshwater systems. *Can. J. Fish. Aquatic Sci.* 58: 870-878.

- Latimer, J.S. and J.G. Quinn. 1996. Historical trends and current inputs of HOCs in an urban estuary: The sedimentary record. *Environ. Sci. Technol.* 30: 623-633.
- Leavitt, P.R., C.S. Brock, C. Ebel and A. Patoine. 2006. Landscape-scale effects of urban nitrogen on a chain of freshwater lakes in central North America. *Limnol. Oceanogr.* 51(5): 2262–2277.
- Lebo, M.E. 1991. Particle-bound phosphorus along an urbanized coastal plain estuary. *Mar. Chem.* 34: 201-216.
- LeMay, L.E. 2007. The impact of drainage ditches on salt marsh flow patterns, sedimentation and morphology: Rowley River, Massachusetts. M.S. Thesis, Virginia Institute of Marine Science, Williamsburg, Virginia, 239 p.
- Litke, D.W. 1999. *Review of Phosphorus Control Measures in the United States and Their Effects on Water Quality*. U.S. Geological Survey, Water Sources Investigations Report 99-4007, National Water Quality Assessment Program, Denver, CO.
- McClelland, J.W., I. Valiela and R.H. Michener. 1997. Nitrogen-stable isotope signatures in estuarine food webs: A record of increasing urbanization in coastal watershed. *Limnol. Oceanogr.* 42: 930-937.
- Meybeck, M. 1982. Carbon, nitrogen and phosphorus transport by world rivers. *Amer. Jour. Sci.* 282: 401-450.
- Meyers, P.A., 1994. Preservation of elemental and isotopic source identification of sedimentary organic matter. *Chem. Geol.* 144: 289-302.
- Mitch, W.J. and J.G. Gosselink. 1993. *Wetlands*, 2nd Edition. Van Nostrand Reinhold, NY.
- Officer, C.B., D.R. Lynch, G.H. Setlock and G.R. Helz. 1984. Recent sedimentation rates in Chesapeake Bay. In: Kennedy, V.S. (ed.). *The Estuary as a Filter*. Academic Press, New York, pp 131-157.
- O'Leary, M.H., 1988. Carbon isotopes in photosynthesis. *Bioscience*, 38:328-336.
- Olsen, C.R., I.L. Larsen, P.D. Lowry, N.H. Cutshall, J.F. Todd and G.T.F. Wong. 1985. Atmospheric fluxes and marsh-soil inventories of ⁷Be and ²¹⁰Pb. *Journal of Geophysical Research –Atmospheres*. 90: 10487-10495.
- Olsen, C.R., H.J. Simpson, T.H. Peng, R.F. Bopp and R.M. Trier. 1981. Sediment mixing and accumulation effects on radionuclide depth profiles in Hudson estuary sediments. *Journal of Geophysical Research*. 86: 11020-11028.

- Orson, R.A., R.L. Simpson and R.E. Good. 1992. A mechanism for accumulation and retention of heavy metals in tidal freshwater marshes of the upper Delaware River Estuary. *Estuar. Coast. Shelf Sci.* 34: 171-186.
- Orson, R.A., R.L. Simpson and R.E. Good. 1990. Rates of sediment accumulation in a tidal freshwater marsh. *J. Sed. Petrology.* 60: 859-869.
- Owens, M. and J.C. Cornwell. 1995. Sedimentary evidence for decreased heavy metal inputs to the Chesapeake Bay. *Ambio* 24: 24-27.
- Perga, M.E. and D. Gerdeaux. 2004. Changes in the $\delta^{13}\text{C}$ of pelagic food webs: The influence of lake area and trophic status on the isotopic signature of whitefish (*Coregonus lavaretus*). *Can. J. Fish. Aquat. Sci.* 61(8): 1485-1492.
- Ponader, K.C., D.F. Charles, T.J. Belton and D.M. Winter. 2008. Total phosphorus inference models and indices for coastal plain streams based on benthic diatom assemblages from artificial substrates. *Hydrobiologia* 610: 139-152.
- Porter, S.D. 2008. *Algal attributes: An autecological classification of algal taxa collected by the National Water-Quality Assessment Program*: U.S. Geological Survey Data Series 329, 18 p.
- Porter, S.D., D.K. Mueller, N.E. Spahr, M.D. Munn and N.M. Dubrovsky. 2008. Efficacy of algal metrics for assessing nutrient and organic enrichment in flowing waters: *Freshwater Biology.* 53: 1036-1054.
- Potapova, M. and D.F. Charles. 2007. Diatom metrics for monitoring eutrophication in rivers of the United States. *Ecological Indicators.* 7: 48-70.
- Potapova, M., D.F. Charles, K.C. Ponader and D.M. Winter. 2004. Quantifying species indicator values for trophic diatom indices: a comparison of approaches. *Hydrobiologia.* 517: 25-41.
- Prahl, F.G., J.R. Ertel, M.A. Goni, M.A. Sparrow and B. Eversmeyer. 1994. Terrestrial organic carbon contributions to sediments on the Washington margin. *Geochim. Cosmochim. Acta.* 58:3035-3048.
- Ridgeway, J. and G. Shimmield. 2002. Estuaries as repositories of historical contamination and their impact on shelf seas. *Estuar. Coast. Shelf Sci.* 55: 903-928.
- Rippey, B. and H. J. Anderson. 1996. Reconstruction of lake P loadings and dynamics using sedimentary record. *Environ. Sci. Technol.* 30: 1786-1789.
- Ritchie, J.C. and McHenry, R.J. 1990. Application of radioactive fallout Cesium-137 for measuring soil erosion and sediment accumulation rates and patterns: A review. *Journal of Environmental Quality.* 19: 215-233.

- Robbins, J.A. 1978. Geochemical and geophysical application of radioactive lead. In: J.O. Nriagu (ed.). *The biogeochemistry of lead in the environment*. Elsevier, Amsterdam.
- Ruttenberg, K.C. 1992. Development of a sequential extraction method for different forms of phosphorus in marine sediments. *Limnol. Oceanogr.* 37: 1460-1482.
- Santschi, P.H., B.J. Presley, T.L. Wade, B. Garcia-Romero and M. Baskaran. 2001. Historical contamination of PAHs, PCBs, DDTs, and heavy metals in Mississippi River Delta, Galveston Bay and Tampa Bay sediment cores. *Mar. Environ. Res.* 52: 51-79.
- Schelske, C.L. and D.A. Hodell. 1995. Using carbon isotopes of bulk sedimentary organic matter to reconstruct the history of nutrient loading and eutrophication in Lake Erie. *Limnol. Oceanogr.* 40: 918-929.
- Schelske, C.L. and D.A. Hodell. 1991. Recent changes in productivity and climate of Lake Ontario detected by isotopic analysis of sediments. *Limnol. Oceanogr.* 36: 961-975.
- Schubel, J. R. and D. J. Hirschberg. 1977. ^{210}Pb determined sedimentation rate, and accumulation of metals in sediments at a station in Chesapeake Bay. *Chesapeake Sci.* 18: 379-382.
- Simpson, R.L., R.E. Good, R. Walker and B.R. Frasco. 1983. The role of Delaware River freshwater tidal wetlands in the retention of nutrients and heavy metals. *J. Environ. Qual.* 12: 41-48.
- Smith, B.N. and S. Epstein. 1971. Two categories of $^{13}\text{C}/^{12}\text{C}$ ratios for higher plants. *Plant Physiol.* 47:380-384.
- Smol, J.P. 2008. *Pollution of Lakes and Rivers: A Paleoenvironmental Perspective*- 2nd edition. Blackwell Publishing, Oxford.
- Sommerfield, C.K. and D.J. Velinsky. 2010. Sediment Accumulation and Marsh Accretion in Tidal Wetlands of a Coastal Plain Estuary. Presented at the 2010 American Geophysical Ocean Sciences Meeting, Portland, OR.
- Ulseth, A.J. and A.E. Hershey. 2005. Natural abundances of stable isotopes trace anthropogenic N and C in an urban stream. *JNABS*: 24(2): 270-289.
- USEPA 2005. Letter to Mr. Kevin Donnelly from US EPA (August 25, 2005) from Mr. Jon Capacasa, Director, WPD. Decision Rationale: TMDL Analysis for the Murderkill River Watershed, US EPA Region III, Philadelphia, PA
- US EPA 1997. Methods for the Determination of Chemical Substances in Marine and Estuarine Environmental Matrices. 2nd edition. September 1997. (NSCEP or NTIS / PB97-127326)

- Ullman, W.J., M. Dean, J.R. Scudlark, J. Newton, H. Mirsajadi, A.K. Aufenkampe, R.L. Hays, B. Dzwonkowski, K.C. Wong, T.E. McKenna, and A.S. Andres. 2010. Contrasting loads of nitrogen and phosphorus to and from the Murderkill Watershed and Estuary, Delaware, USA. (11 January 2010) OOB Discussion.
- Ulseth, A.J. and A.E. Hershey. 2005. Natural abundances of stable isotopes trace anthropogenic N and C in an urban stream. *JNABS*: 24(2): 270-289.
- Valette-Silver, N.J. 1993. The use of sediment cores to reconstruct historical trends in contamination of estuarine and coastal sediments. *Estuaries*. 16: 577-588.
- Van Breemen et al. 2002. Where did all the nitrogen go? Fate of nitrogen inputs to large watersheds in the NE USA. *Biogeochemistry* 57/58: 267-293.
- van Dam, H., A. Mertens, and J. Sinkeldam. 1994. A coded checklist and ecological indicator values of freshwater diatoms from The Netherlands. *Neth. J. Aquat. Ecol.* 28: 117-133.
- Van Metre, P.C. and Callender, E. 1997. Identifying water-quality trends from ^{210}Pb using dated sediment cores from White Rock Lake Reservoir, Dallas, Texas. *Journal of Paleolimnology*. 17: 239-249.
- Velinsky, D.J., Riedel, G.R., J.T.F. Ashley and J.Cornwell. 2010. River sediment investigation near Poplar Point in the tidal Anacostia River, Washington, DC. *Environ. Monior. Assess* (submitted).
- Velinsky, D.J., C. Sommerfield and D. Charles. 2010. Vertical profiles of radioisotopes, contaminants, nutrients and diatoms in sediment cores from the tidal Christina River Basin: A historical analysis. Report submitted to Dr. R. Greene (DNREC; Division of Water Resources; State of Delaware, Dover DE). Patrick Center Report 09-02, The Academy of Natural Sciences, Philadelphia, PA.
- Velinsky, D.J., D.F. Charles and J. Ashley. 2007. Contaminant Sediment Profiles of the St. Jones River Marsh, Delaware: A Historical Analysis. Final Report submitted to DNREC. PCER Report No. 07-05. The Academy of Natural Sciences, Patrick Center for Environmental Research, Philadelphia, PA.
- Wallbrink, P.J., D.E. Walling and Q. He. 2002. Radionuclide measurement using HPGe gamma spectrometry. In: F. Zapata (Editor), *Handbook for the Assessment of Soil Erosion and Sedimentation Using Environmental Radionuclides*. Kluwer Academic Publishers, Dordrecht, pp. 67-96.
- Zhang, X., E.R. Christensen and M.F. Gin. 1993. PAHs in dated sediment from Green Bay and Lake Michigan. *Estuaries* 16: 638-652.
- Zimmerman, A.R. and E.A. Canuel. 2002. Sediment geochemical records eutrophication in the mesohaline Chesapeake Bay. *Limnol. Oceanogr.* 47: 1084-1093.

Tables

Table 1. Core locations and collection dates.

Name	Abbreviation	Date	Lat (N)	Long (W)	Depth of Core (cm)
Downstream of Bay Rd	MK-1A	10-Oct-08	39° 00.587'	75° 26.963'	73.8
	MK-1B	10-Oct-08	39° 00.584'	75° 26.962'	82.0
Near Outfall Creek	MK-2A	10-Oct-08	39° 00.662'	75° 26.390'	91.0
	MK-2B	10-Oct-08	39° 00.662'	75° 26.391'	94.0
Mid River	MK-3A	10-Oct-08	39° 01.556'	75° 24.751'	98.5
	MK-3B	10-Oct-08	39° 01.555'	75° 24.750'	98.0
Near USGS Gage	MK-4A	10-Oct-08	39° 02.997'	75° 23.523'	104.0
	MK-4B	10-Oct-08	39° 02.998'	75° 23.521'	96.0

Stations are listed from upstream to downstream.

Table 2. Bulk sediment properties and radioisotope data for core MK-1.

Chem ID	Interval	Water Fraction	Dry Bulk Density (g/cm ³)	Accumulated Mass (g/cm ²)	LOI (%)	Total ²¹⁰ Pb (dpm/g)	Excess ²¹⁰ Pb (dpm/g)	¹³⁷ Cs (dpm/g)	z (cm)	Age Model*
	0-2	0.74	0.314	1.29	26.8	8.33	7.71	0.46	2	2005
	2-4	0.73	0.329		26.4				4	2002
	4-6	0.72	0.336	1.30	27.3	7.58	6.85	0.43	6	2000
	6-8	0.74	0.315		28.8				8	1997
	8-10	0.71	0.354	1.51	24.6	8.38	7.54	0.37	10	1994
	10-12	0.68	0.399		21.9				12	1991
	12-14	0.69	0.389	0.78	21.0	7.49	6.55	0.42	14	1989
	14-16	0.68	0.398	0.80	23.1	8.19	6.82	0.52	16	1986
	16-18	0.73	0.328	1.24	23.9	6.91	5.85	0.54	18	1983
	18-20	0.75	0.290		25.7				20	1980
	20-22	0.78	0.253	1.00	27.6	5.23	4.09	1.34	22	1978
	22-24	0.79	0.247		28.6				24	1975
	24-26	0.82	0.197	0.82	33.9	4.30	3.33	1.73	26	1972
	26-28	0.81	0.211		31.1				28	1969
	28-30	0.81	0.217	0.97	29.5	4.16	3.32	2.27	30	1967
	30-32	0.77	0.266		26.4				32	1964
	32-34	0.77	0.271	1.14	26.8	3.05	2.10	0.84	34	1961
	34-36	0.75	0.300		25.8				36	1958
	36-38	0.75	0.297	1.10	26.4	2.15	1.20	0.25	38	1956
	38-40	0.78	0.252		31.3				40	1953
	40-42	0.76	0.280	1.08	28.3	2.30	1.29	0.10	42	1950
	42-44	0.78	0.258		29.6				44	1947
	44-46	0.83	0.187	0.69	39.2	2.68	2.00	0.05	46	1945
	46-48	0.86	0.158		49.7				48	1942
	48-50	0.86	0.152	0.59	56.3	1.30	0.60	-0.04	50	1939
	50-52	0.87	0.145		58.8				52	1936
	52-54	0.86	0.151	0.62	59.7	1.24	1.00	0.06	54	1934
	54-56	0.85	0.161		58.7				56	1931
	56-58	0.86	0.155	0.63	70.5	0.86	0.62	-0.08	58	1928
	58-60	0.86	0.159		60.8				60	1925
	60-62	0.85	0.160		60.6					
	62-64	0.84	0.178		55.9					
	64-66	0.85	0.172		56.5					
	66-68	0.83	0.187		55.1					
	68-70	0.83	0.187		54.6					
	70-72	0.81	0.217		53.8					
	72-74	0.83	0.186		53.0					

74-76	0.82	0.197	51.0
76-78	0.83	0.194	49.8
78-80	0.82	0.198	51.6
80-82	0.83	0.191	49.7

*Based on a ^{210}Pb sedimentation rate of 0.74 cm/yr. *Note: Cells for accumulated mass, total ^{210}Pb , excess ^{210}Pb and ^{137}Cs have been merged in cases where adjacent samples were physically added together for measuring radionuclide activities. For example, MK-1 samples 0-2cm and 2-4cm were dried and combusted separately but combined for radionuclide analysis.

Table 3. Bulk sediment properties and radioisotope data for MK-2.

Chem ID	Interval	Water Fraction	Dry Bulk Density (g/cm ³)	Accumulated Mass (g/cm ²)	LOI (%)	Total ²¹⁰ Pb (dpm/g)	Excess ²¹⁰ Pb (dpm/g)	¹³⁷ Cs (dpm/g)	z (cm)	Age Model*
	0-2	0.84	0.180	0.69	51.3	10.91	10.56	0.09	2	2005
	2-4	0.85	0.166		59.3				4	2002
	4-6	0.86	0.159	0.65	61.9	11.68	11.13	0.11	6	2000
	6-8	0.85	0.163		55.2				8	1997
	8-10	0.85	0.163	0.72	49.7	8.97	8.87	0.26	10	1994
	10-12	0.83	0.196		41.9				12	1991
	12-14	0.86	0.154	0.56	54.6	8.57	8.27	0.01	14	1989
	14-16	0.88	0.128		56.7				16	1986
	16-18	0.89	0.119	0.47	52.3	6.49	6.79	0.40	18	1983
	18-20	0.89	0.114		53.2				20	1980
	20-22	0.89	0.115	0.55	46.9	6.11	5.74	0.58	22	1978
	22-24	0.85	0.161		34.8				24	1975
	24-26	0.84	0.178	0.80	31.9	4.63	3.90	1.78	26	1972
	26-28	0.81	0.220		29.0				28	1969
	28-30	0.79	0.237	0.99	30.8	3.91	3.22	4.43	30	1967
	30-32	0.78	0.259		29.0				32	1964
	32-34	0.77	0.265	1.07	27.6	2.54	1.97	2.08	34	1961
	34-36	0.77	0.272		25.7				36	1958
	36-38	0.78	0.256	1.03	29.1	2.60	1.89	0.70	38	1956
	38-40	0.78	0.258		29.1				40	1953
	40-42	0.78	0.259	1.03	30.5	2.40	1.61	0.32	42	1950
	42-44	0.78	0.253		32.0				44	1947
	44-46	0.82	0.206	0.77	45.3	1.52	1.02	0.11	46	1945
	46-48	0.84	0.181		55.6				48	1942
	48-50	0.84	0.177	0.69	59.3	0.84	1.04	-0.10	50	1939
	50-52	0.85	0.170		61.6				52	1936
	52-54	0.85	0.170	0.67	61.2	0.71	0.36	0.23	54	1934
	54-56	0.85	0.162		65.8				56	1931
	56-58	0.85	0.160	0.64	68.2	0.53	-0.06	-0.04	58	1928
	58-60	0.86	0.158		68.2				60	1925
	60-62	0.86	0.155	0.61	71.3	0.65	0.73	-0.08	62	1922
	62-64	0.86	0.152		71.1				64	1920
	64-66	0.86	0.154	0.60	70.6	-0.05	-0.13	0.01	66	1917
	66-68	0.87	0.145		70.7				68	1914
	68-70	0.87	0.147	0.56	69.4	1.04	1.24	-0.10	70	1911
	70-72	0.88	0.135		67.0				72	1909
	72-74	0.88	0.124	0.51	64.6	-0.07	-0.38	0.01	74	1906

74-76	0.88	0.133		55.4				76	1903
76-78	0.87	0.143	0.57	51.9	0.88	0.14	-0.10	78	1900
78-80	0.87	0.142		51.4				80	1898
80-82	0.87	0.143	0.59	51.8	0.62	0.68	-0.07	82	1895
82-84	0.86	0.151		53.6				84	1892
84-86	0.86	0.157	0.63	53.1	0.55	0.43	0.06	86	1889
86-88	0.85	0.160		53.7				88	1887
88-90	0.85	0.169	0.68	51.4	0.77	0.41	0.02	90	1884
90-92	0.85	0.169		52.2				92	1881
92-94	0.84	0.182	0.73	49.3	0.07	-0.24	0.16	94	1878

*Based on ²¹⁰Pb sedimentation rate of 0.74 cm/yr

Table 4. Bulk sediment properties and radioisotope data for MK-3.

Chem ID	Interval	Water Fraction	Dry Bulk Density (g/cm ³)	Accumulated Mass (g/cm ²)	LOI (%)	Total ²¹⁰ Pb (dpm/g)	Excess ²¹⁰ Pb (dpm/g)	¹³⁷ Cs (dpm/g)	z (cm)	Age Model*
	0-2	0.74	0.304	1.12	33.9	11.85	11.44	0.29	2	2004
	2-4	0.78	0.256		46.1				4	2000
	4-6	0.73	0.321	0.64	40.3	8.67	7.51	0.23	6	1996
	6-8	0.77	0.270	0.54	44.0	7.11	6.73	0.46	8	1993
	8-10	0.82	0.203	0.84	52.0	6.18	5.62	1.12	10	1989
	10-12	0.81	0.218		48.3				12	1985
	12-14	0.79	0.239	0.88	38.3	6.33	5.75	3.61	14	1981
	14-16	0.82	0.203		35.1				16	1977
	16-18	0.83	0.191	0.78	33.5	4.22	3.74	4.70	18	1973
	18-20	0.82	0.197		35.3				20	1970
	20-22	0.83	0.186	0.71	37.9	4.27	4.24	2.04	22	1966
	22-24	0.85	0.168		37.5				24	1962
	24-26	0.86	0.158	0.65	36.3	3.92	3.44	0.21	26	1958
	26-28	0.85	0.169		34.4				28	1954
	28-30	0.81	0.220	0.93	32.1	2.44	1.68	0.21	30	1950
	30-32	0.79	0.247		27.3				32	1946
	32-34	0.81	0.221	0.93	32.0	2.40	1.43	0.08	34	1943
	34-36	0.79	0.244		29.5				36	1939
	36-38	0.81	0.216	0.82	26.4	1.36	0.72	-0.12	38	1935
	38-40	0.83	0.196		34.3				40	1931
	40-42	0.85	0.168	0.70	33.9	1.23	1.03	-0.14	42	1927
	42-44	0.84	0.181		38.1				44	1923
	44-46	0.77	0.269	1.07	29.1	1.13	0.58	-0.04	46	1920
	46-48	0.77	0.267		37.0				48	1916
	48-50	0.77	0.262	0.97	27.2	1.60	0.77	-0.08	50	1912
	50-52	0.80	0.225		30.4				52	1908
	52-54	0.83	0.196		29.2					
	54-56	0.80	0.223		28.4					
	56-58	0.74	0.311		24.6					
	58-60	0.68	0.400		25.0					
	60-62	0.68	0.397		25.0					
	62-64	0.72	0.343		30.7					
	64-66	0.75	0.290		35.6					
	66-68	0.75	0.292		40.7					
	68-70	0.75	0.295		37.9					
	70-72	0.74	0.311		37.5					
	72-74	0.75	0.296		36.1					

74-76	0.76	0.278	39.3
76-78	0.77	0.273	38.1
78-80	0.78	0.256	37.8
80-82	0.77	0.266	35.5
82-84	0.76	0.288	30.8
84-86	0.68	0.401	21.2
86-88	0.67	0.415	17.7
88-90	0.64	0.460	16.6
90-92	0.60	0.525	13.1
92-94	0.57	0.588	12.0
94-96	0.58	0.575	12.0
96-98	0.54	0.635	11.1

*Based on a ²¹⁰Pb sedimentation rate of 0.60 cm/yr

Table 5. Bulk sediment properties and radioisotope data for MK-4.

Chem ID	Interval	Water Fraction	Dry Bulk Density (g/cm ³)	Accumulated Mass (g/cm ²)	LOI (%)	Total ²¹⁰ Pb (dpm/g)	Excess ²¹⁰ Pb (dpm/g)	¹³⁷ Cs (dpm/g)	z (cm)	Age Model*
	0-2	0.58	0.570	1.14	27.0	9.93	8.84	0.32	2	2002
	2-4	0.60	0.535	1.07	17.0	10.08	9.02	0.17	4	1996
	4-6	0.64	0.466	0.93	37.8	8.46	7.39	0.28	6	1989
	6-8	0.63	0.479	0.96	22.4	6.25	5.45	0.85	8	1983
	8-10	0.66	0.436	1.81	28.4	4.97	4.28	2.00	10	1977
	10-12	0.64	0.467		20.2				12	1971
	12-14	0.69	0.383	0.77	25.5	3.79	3.04	2.06	14	1964
	14-16	0.71	0.348	0.70	29.0	4.03	3.21	1.80	16	1958
	16-18	0.71	0.347	1.43	26.7	2.89	2.12	0.63	18	1952
	18-20	0.70	0.366		24.7				20	1946
	20-22	0.66	0.434	0.87	18.4	2.27	1.17	0.24	22	1939
	22-24	0.69	0.380	0.76	20.0	2.10	1.18	0.00	24	1933
	24-26	0.67	0.408	0.82	17.1	1.92	0.66	0.01	26	1927
	26-28	0.66	0.439	0.88	15.0	1.91	0.77	0.00	28	1921
	28-30	0.65	0.451	0.90	14.2	1.46	0.43	-0.02	30	1914
	30-32	0.65	0.455	0.91	13.7	2.24	1.22	0.04	32	1908
	32-34	0.60	0.537	1.07	12.2	1.49	0.17	0.01	34	1902
	34-36	0.58	0.566	1.13	11.7	1.31	0.31	-0.04	36	1896
	36-38	0.56	0.608	1.22	12.1	1.13	-0.23	0.03	38	1889
	38-40	0.55	0.629	1.26	11.8	1.77	0.76	-0.04	40	1883
	40-42	0.54	0.646	1.29	11.7	1.22	0.05	0.11	42	1877
	42-44	0.52	0.691	1.38	12.1	1.28	-0.10	-0.04	44	1871
	44-46	0.54	0.638	1.28	11.5	0.63	-0.51	-0.04	46	1864
	46-48	0.52	0.682	1.36	11.6	1.76	0.44	0.03	48	1858
	48-50	0.50	0.728	1.46	9.7	1.70	0.36	-0.03	50	1852
	50-52	0.48	0.763		9.6					
	52-54	0.52	0.683		10.5					
	54-56	0.57	0.592		12.8					
	56-58	0.59	0.554		12.4					
	58-60	0.60	0.526		13.9					
	60-62	0.61	0.517		11.7					
	62-64	0.61	0.514		13.4					
	64-66	0.64	0.458		13.1					
	66-68	0.62	0.503		13.0					
	68-70	0.62	0.490		12.0					
	70-72	0.62	0.493		13.0					
	72-74	0.61	0.508		11.5					

74-76	0.59	0.547	13.0
76-78	0.58	0.566	11.7
78-80	0.59	0.555	13.9
80-82	0.62	0.502	14.0
82-84	0.57	0.583	14.0
84-86	0.60	0.539	13.6
86-88	0.62	0.494	18.5
88-90	0.65	0.444	19.2
90-92	0.64	0.468	19.1
92-94	0.62	0.500	15.2
94-96	0.62	0.495	17.6

*Based on a ^{210}Pb sedimentation rate of 0.33 cm/yr

Table 6. Summary data for radioisotope analysis and dating.

Location	Core	Radioisotope Inventory		Accretion Rate		Mass Accumulation Rate		Focusing Factor	
		xsPb-210 dpm/cm ²	Cs-137 dpm/cm ²	Pb-210 cm/yr	Cs-137 cm/yr	xsPb-210 g/cm ² -yr	Cs-137 g/cm ² -yr	xsPb-210	Cs-137
Upstream	MK-1	65.80	9.38	0.74	0.71	0.20	0.19	2.35	0.45
	MK-2	47.29	10.04	0.74	0.71	0.13	0.12	1.69	0.48
	MK-3	44.81	10.03	0.60	0.44	0.14	0.10	1.60	0.48
Downstream	MK-4	53.40	9.20	0.33	0.31	0.17	0.16	1.91	0.44

Table 7. Concentrations of various parameters for Core MK-1.

CHEM ID	Interval cm	Mid-Point cm	Solids %	Age Model yr	$\delta^{13}\text{C}$ ‰	$\delta^{15}\text{N}$ ‰	TC %	TN %	TSP %	C/N Atomic
2028	0-2	1	27.1	2006	-17.12	9.96	11.56	0.73	0.107	18.5
2030	4-6	5	28.5	2000	-17.30	9.45	10.77	0.68	0.065	18.4
2032	8-10	9	30.1	1995	-18.27	8.68	9.63	0.65	0.071	17.3
2033	10-12	11	31.7	1992	-17.53	8.63	9.18	0.62	0.075	17.2
2034	12-14	13	30.2	1990	-17.67	8.96	8.55	0.65	0.076	15.4
2035	14-16	15	26.2	1987	-17.72	9.97	9.45	0.72	0.082	15.3
2036	16-18	17	27.0	1984	-17.73	8.68	10.14	0.74	0.084	16.1
2037	18-20	19	22.1	1981	-17.58	8.84	11.59	0.74	0.077	18.4
2038	20-22	21	25.0	1979	-17.46	8.67	9.73	0.61	0.075	18.6
2039	22-24	23	23.1	1976	-17.73	7.97	10.19	0.65	0.074	18.4
2042	28-30	29	21.5	1968	-19.14	5.74	11.53	0.72	0.065	18.6
2044	32-34	33	48.1	1963	-21.14	4.88	11.90	0.66	0.065	21.2
2046	36-38	37	23.8	1957	-22.84	4.70	11.19	0.60	0.062	21.7
2048	40-42	41	24.0	1952	-26.36	3.74	11.84	0.67	0.065	20.5
2050	44-46	45	14.7	1946	-26.52	1.53	25.36	1.23	0.086	24.0
2054	52-54	53	14.4	1936	-27.12	0.59	31.18	1.54	0.080	23.6
2059	62-64	63	16.7	1922	-26.97	0.58	27.36	1.44	0.071	22.2
2064	72-74	73	19.1	1909	-25.72	0.83	25.99	1.25	0.072	24.3
2067	78-80	79	19.5	1900	-25.52	1.15	23.88	1.18	0.066	23.6
Spartina alt.	Surface		NA	NA	-13.61	9.02	10.36	1.55	NA	30.4

Table 8. Concentrations of various parameters for Core MK-2.

CHEM ID	Interval cm	Mid-Point cm	Solids %	Age Model yr	$\delta^{13}\text{C}$ ‰	$\delta^{15}\text{N}$ ‰	TC %	TN %	TSP %	C/N Atomic
2069	0-2	1	16.3	2006	-15.08	6.98	27.72	1.28	0.119	25.2
2071	4-6	5	16.8	2000	-16.55	6.67	25.17	1.53	0.107	19.2
2073	8-10	9	19.2	1995	-15.80	7.48	17.29	0.88	0.072	22.8
2075	12-14	13	14.2	1990	-14.84	7.29	26.33	1.31	0.080	23.4
2077	16-18	17	12.2	1984	-14.87	6.34	21.35	0.94	0.063	26.5
2079	20-22	21	12.4	1979	-15.09	6.71	21.12	0.95	0.059	26.0
2081	24-26	25	17.4	1973	-15.97	5.57	12.73	0.64	0.061	23.0
2083	28-30	29	20.6	1968	-16.08	5.21	12.63	0.58	0.057	25.4
2085	32-34	33	22.2	1963	-18.96	4.20	11.73	0.58	0.058	23.4
2087	36-38	37	21.8	1957	-21.71	3.00	12.40	0.65	0.058	22.1
2090	42-44	43	18.8	1949	-26.51	0.56	25.70	1.22	0.065	24.5
2094	50-52	51	15.1	1938	-25.84	0.03	32.14	1.69	0.079	22.2
2099	60-62	61	15.4	1925	-24.72	-0.12	34.33	1.61	0.065	24.8
2106	74-76	75	12.3	1906	-23.35	0.13	28.39	1.29	0.070	25.6
2113	88-90	89	16.6	1887	-23.35	0.55	23.20	1.05	0.064	25.8
Spartina alt.	Surface		NA	NA	-12.71	3.79	48.53	0.79	NA	71.7

Concentrations on a dry weight basis. NA- Not analyzed.

Table 9. Concentrations of various parameters for Core MK-3.

CHEM ID	Interval cm	Mid-Point cm	Solids %	Age Model yr	$\delta^{13}\text{C}$ ‰	$\delta^{15}\text{N}$ ‰	TC %	TN %	TSP %	C/N Atomic
2116	0-2	1	26.51	2005	-15.93	1.15	12.69	0.69	0.087	21.5
2118	4-6	5	29.43	1999	-15.12	0.94	14.73	0.53	0.058	32.4
2120	8-10	9	21.25	1992	-14.19	1.41	22.21	0.70	0.045	36.8
2122	12-14	13	22.88	1985	-14.21	0.03	15.59	0.63	0.071	28.9
2123	14-16	15	18.10	1982	-14.14	0.81	16.66	0.63	0.060	31.0
2124	16-18	17	18.34	1979	-14.47	0.15	14.88	0.62	0.069	27.9
2125	18-20	19	16.32	1975	-13.99	0.61	16.48	0.61	0.054	31.8
2126	20-22	21	16.47	1972	-14.21	0.34	16.12	0.60	0.054	31.4
2128	24-26	25	13.35	1965	-13.14	0.68	24.41	0.71	0.051	40.3
2130	28-30	29	19.58	1959	-14.62	0.23	12.10	0.56	0.120	25.3
2132	32-34	33	20.35	1952	-14.41	0.06	12.80	0.48	0.049	31.4
2134	36-38	37	20.68	1945	-14.85	0.38	10.93	0.43	0.054	29.7
2136	40-42	41	14.84	1939	-15.42	0.41	15.05	0.62	0.043	28.2
2140	48-50	49	22.94	1925	-14.99	0.65	10.54	0.44	0.051	28.2
2145	58-60	59	31.26	1909	-14.70	0.39	10.81	0.48	0.044	26.2
2151	70-72	71	25.97	1889	-16.20	0.00	3.36	0.21	0.058	19.0
2161	90-92	91	46.84	1855	-19.69	0.02	15.62	0.85	0.042	21.4
Spartina alt.	Surface		NA	NA	-12.74	3.74	46.57	0.71	NZ	76.5

Concentrations on a dry weight basis. NA- Not analyzed.

Table 10. Concentrations of various parameters for Core MK-4.

CHEM ID	Interval	Mid-Point	Solids	Age Model	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	TC	TN	TSP	C/N
	cm	cm	%	yr	‰	‰	%	%	%	Atomic
2165	0-2	1	44.1	2003	-15.28	3.34	14.12	0.54	0.157	30.4
2167	4-6	5	37.9	1991	-16.08	4.65	7.37	0.48	0.205	18.1
2169	8-10	9	38.3	1979	-16.41	4.65	6.77	0.43	0.199	18.6
2170	10-12	11	31.4	1973	-16.77	6.11	5.34	0.38	0.069	16.5
2171	12-14	13	34.9	1967	-16.14	3.75	7.05	0.45	0.069	18.3
2172	14-16	15	33.2	1961	-15.81	3.22	8.48	0.42	0.063	23.4
2173	16-18	17	30.8	1954	-15.75	2.61	8.06	0.48	0.070	19.5
2174	18-20	19	30.4	1948	-15.69	2.24	9.00	0.49	0.059	21.5
2175	20-22	21	39.2	1942	-15.56	2.43	5.55	0.29	0.054	22.3
2176	22-24	23	36.5	1936	-15.95	1.99	5.91	0.33	0.058	21.2
2179	28-30	29	32.9	1918	-15.83	1.86	6.25	0.33	0.049	22.3
2181	32-34	33	45.4	1906	-16.66	2.82	3.79	0.28	0.050	15.8
2183	36-38	37	45.5	1894	-15.72	3.10	3.73	0.29	0.048	14.9
2185	40-42	41	46.3	1882	-15.04	2.66	4.41	0.31	0.048	16.5
2187	44-46	45	48.1	1870	-15.91	3.06	2.74	0.22	0.052	14.6
2189	48-50	49	49.8	1857	-16.64	3.74	2.32	0.21	0.049	13.0
2191	52-54	53	51.6	1845	-15.78	3.21	3.12	0.24	0.049	15.3
2193	56-58	57	40.5	1833	-14.94	2.03	5.35	0.33	0.053	18.8
2199	68-70	69	40.5	1797	-16.01	1.79	4.17	0.25	0.046	19.5
2203	76-78	77	41.4	1773	-15.75	0.61	4.52	0.31	0.046	17.3
2206	82-84	83	41.1	1754	-15.65	0.53	5.35	0.34	0.047	18.2
2209	88-90	89	37.5	1736	-14.56	0.86	7.71	0.44	0.051	20.2
2212	94-96	95	39.2	1718	-14.24	1.64	5.45	0.31	0.048	20.6
Spartina alt.	Surface				-13.23	4.17	45.23	0.70		75.4

Concentrations on a dry weight basis. NA- Not analyzed.

Table 11. Diatom Metrics determined from species identification: MK-1.

Core ID/ Depth Interval (cm)	Mid-Depth (cm)	Eutrophic	Meso/ eutrophic	Mesotrophic	Oligotrophic	Oligo/ mesotrophic	Unknow
0-2	1	38.0	3.3	0.0	0.0	0.0	57.8
4-6	5	48.0	1.8	0.5	0.3	0.0	49.0
8-10	9	45.5	1.3	0.0	0.0	0.0	50.8
12-14	13	44.0	1.5	0.0	0.8	0.8	51.3
16-18	17	39.5	4.5	0.0	0.0	0.0	55.0
20-22	21	37.5	7.8	0.0	0.0	0.0	53.5
22-24	23	37.3	10.3	0.0	1.3	0.0	51.3
28-30	29	54.3	4.0	0.8	0.3	0.0	40.8
32-34	33	61.3	7.8	0.0	2.0	0.0	29.0
36-38	37	68.3	5.3	0.0	0.8	0.0	24.8
44-46	45	12.8	9.0	2.0	51.8	1.0	22.5
52-54	53	9.3	7.3	0.8	70.3	0.5	12.0
62-64	63	18.8	11.5	4.3	31.5	5.0	29.0
72-74	73	24.0	10.5	8.5	20.5	6.5	29.3
78-80	79	22.5	9.8	7.5	9.8	8.0	39.8

Metrics based on van Dam et al. (1994)

Table 12. Diatom Metrics determined from species identification: MK-2.

Core ID/ Depth Interval (cm)	Mid-Depth (cm)	Eutrophentic	Meso/ eutrophentic	Mesotrophentic	Oligotrophentic	Oligo/ mesotrophentic	Unknow
0-2	1	21.8	4.3	2.5	0.0	0.0	70.3
4-6	5	20.3	14.0	1.0	3.8	0.0	61.0
8-10	9	41.8	4.8	0.5	2.3	0.5	49.8
12-14	13	27.5	16.3	0.0	0.0	0.0	56.3
16-18	17	26.0	29.0	0.3	0.0	0.0	44.8
20-22	21	26.8	12.3	1.0	0.0	0.3	59.3
24-26	25	53.0	3.3	3.5	0.3	0.0	40.0
28-30	29	59.5	1.3	3.0	0.0	0.5	34.5
32-34	33	53.5	1.3	3.0	0.8	2.3	39.3
36-38	37	44.5	6.8	0.5	9.0	0.5	38.8
42-44	43	15.0	7.0	1.5	43.5	1.5	31.5
50-52	53	3.0	12.5	0.3	72.5	0.0	11.8
60-62	63	27.5	2.3	3.5	34.5	1.3	31.0
88-90	89	21.3	7.5	2.5	6.0	1.8	60.8

Metrics based on van Dam et al. (1994)

Table 13. Diatom Metrics determined from species identification: MK-3.

Core ID/ Depth Interval (cm)	Mid-Depth (cm)	Eutrophentic	Meso/ eutrophentic	Mesotrophentic	Oligotrophentic	Oligo/ mesotrophentic	Unknow
0-2	1	37.5	0.3	1.3	0.0	0.0	60.0
4-6	5	44.5	0.5	0.5	0.0	0.0	53.5
8-10	9	65.8	0.5	0.0	0.0	0.0	32.8
12-14	13	62.8	1.8	0.5	0.0	0.0	35.0
14-16	15	58.0	2.8	0.0	0.0	0.5	37.8
16-18	17	51.5	2.8	0.3	0.0	0.8	44.8
20-22	21	37.8	6.3	0.0	0.0	0.0	56.0
24-26	25	35.0	1.8	1.0	0.0	0.0	62.0
28-30	29	24.0	0.8	0.0	0.0	1.8	72.5
32-34	33	35.3	4.3	1.8	0.0	1.5	56.3
36-38	37	23.3	5.3	4.5	0.0	0.0	67.0
40-42	41	30.3	4.3	9.0	0.0	3.5	52.5
48-50	49	22.3	8.8	4.8	0.0	1.3	63.0
58-60	59	25.0	15.5	2.0	0.0	2.3	55.3
70-72	71	26.8	10.3	3.0	2.0	1.5	55.0
90-92	91	39.0	2.0	1.3	0.0	0.5	57.3

Metrics based on van Dam et al. (1994)

Table 14. Diatom Metrics determined from species identification: MK-4.

Core ID/ Depth Interval (cm)	Mid-Depth (cm)	Eutrophentic	Meso/ eutrophentic	Mesotrophentic	Oligotrophentic	Oligo/ mesotrophentic	Unknow
0-2	1	18.0	0.8	0.0	0.0	0.0	80.0
4-6	5	27.6	0.5	0.0	0.3	0.0	70.7
8-10	9	27.9	1.7	1.2	0.5	0.0	68.7
10-12	13	34.0	0.0	0.0	0.3	2.0	63.7
12-14	15	25.7	0.0	0.3	0.3	0.0	73.7
16-18	17	28.0	1.0	0.7	0.0	1.3	69.0
20-22	21	23.8	0.3	0.0	0.0	0.9	74.9
24-26	25	27.4	0.3	0.7	0.3	1.0	70.4
28-30	29	24.7	0.3	0.0	0.0	5.7	69.3
40-42	43	17.7	0.0	1.3	0.0	0.0	81.1
56-58	57	25.5	0.0	0.7	0.0	1.0	72.9

Metrics based on van Dam et al. (1994)

Figures





Figure 1a,b. Generalized schematic of nitrogen and phosphorus cycling in wetlands. SOP – soluble organic phosphorus. In marine sediments, dissolved sulfide (from sulfate reduction) can bind reduced iron, allowing for greater movement of porewater P from the sediments to the overlying water. (Images taken from Mitch and Gosselink, 1993)



Figure 2. Coring locations within the tidal Murderkill River system. Key: The Kent County WWTP is located up the tidal creek near MK-2(A,B).



Figure 3. Tripod and pulley system used to retrieve push-piston cores the Murderkill River. Upper picture is from the upper tidal river (MK-1) while the lower picture is from MK-4, near the tidal inlet and the Delaware Estuary.




Figure 4. Profiles of LOI, dry-bulk density, and excess ^{210}Pb and ^{137}Cs activity for core MK-1B. The regression line used to compute the ^{210}Pb accretion rate is shown.




Figure 5. Profiles of LOI, dry-bulk density, and excess ^{210}Pb and ^{137}Cs activity for core MK-2B. The regression line used to compute the ^{210}Pb accretion rate is shown.




Figure 6. Profiles of LOI, dry-bulk density, and excess ^{210}Pb and ^{137}Cs activity for core MK-3B. The regression line used to compute the ^{210}Pb accretion rate is shown.




Figure 7. Profiles of LOI, dry-bulk density, and excess ^{210}Pb and ^{137}Cs activity for core MK-4B. The regression line used to compute the ^{210}Pb accretion rate is shown.




Figure 8. Sediment organic carbon, total nitrogen, C/N and sediment phosphorus distribution with depth in the marshes of the Murderkill River.




Figure 9. Depth distribution of the isotopic composition of sediment N ($\delta^{15}\text{N}$) and C ($\delta^{13}\text{C}$) from the tidal Murderkill River.




Figure 10. Stratigraphic diagram of diatom species with relative abundances > 5% in at least a sample from core MK-1. Dashed line represents the major shift in diatom assemblages from freshwater *S. exiguiformis* dominant to marine-brackish dominant coastal species.




Figure 11. Stratigraphic diagram of diatom species with relative abundances > 5% in at least a sample from core MK-2. Dashed line represents the major shift in diatom assemblages from freshwater *S. exiguiformis* dominant to marine-brackish dominant coastal species.




Figure 12. Stratigraphic diagram of diatom species with relative abundances > 5% in at least a sample from core MK-3. Dashed line represents the major shift in diatom assemblages from diverse, mixed marine and eutrophic freshwater species to an assemblage with abundant subaerial *Luticola mutica*.




Figure 13. Stratigraphic diagram of diatom species with relative abundances > 5% in at least a sample from core MK-4.

% Diatom Metric (van Dam et al, 1994)




Figure 14. Diatom metrics for the cores of the tidal Christina River using the van Dam et al indices. (1994).




Figure 15. Relationship between total sediment N and P, and the diatom metric for eutrophic species in each core.




Figure 16. Relationship between the accumulation rate of sediment N and P and the diatom metric for eutrophentic species in each core.




Figure 17. Concentrations of total sediment nitrogen (TN) and the nitrogen isotopic composition of TN ($\delta^{15}\text{N-TN}$) from the 1870s to 2008.




Figure 18. Concentrations of total sediment phosphorus (TSP) and the carbon isotopic composition of total carbon ($\delta^{13}\text{C-TC}$) from 1870s to 2008.




Figure 19. Diatom metric for eutrophentic species from 1890s to 2003.





Figure 20. Nitrogen and phosphorus accumulation rates over time in tidal wetlands in the Murderkill River. Linear interpolation was used for sections of sediment that were not analyzed for N or P. The dotted line is at approximately 1975, when the KC WWTP becomes active.

Appendices

Excel File with Data and QA